

This test paper has 4 questions and total of 20 marks. It can not be taken from the examination room. Calculators are allowed. SHOW ALL YOUR WORK. Duration: 50 minutes.

NAME: SOLUTIONS

STUDENT NUMBER:

1. [5 Marks] Let $w = 1 + 4i$ and let $z = -3 + 2i$. Compute the following:

• $w + z$ $(1+4i) + (-3+2i) = (1-3) + (4+2)i = -2 + 6i$

• $z - w$ $(-3+2i) - (1+4i) = (-3-1) + (2-4)i = -4 - 2i$

• wz $(1+4i)(-3+2i) = -3 + 2i - 12i + 8i^2 = -11 - 10i$

• z/w

$$\frac{-3+2i}{1+4i} = \frac{-3+2i}{1+4i} \frac{1-4i}{1-4i}$$

$$= \frac{(-3+2i)(1-4i)}{1+16} = \frac{-3+12i+2i-8i^2}{17}$$

$$= \frac{5}{17} + \frac{14}{17}i$$

2. [5 Marks] Let $p(x)$ be the polynomial $p(x) = x^2 + x + 1$.

- Use the quadratic formula to find the two complex roots of $p(x)$.
- Roughly plot each root of $p(x)$ in the complex plane and express it in polar form (you may use degrees or radians). (Useful fact: $\cos(2\pi/3) = \cos(120^\circ) = -\frac{\sqrt{3}}{2}$.)
- Let z_1 be the root of $p(x)$ with a **positive** imaginary component (i.e. $z_1 = a + bi$, with $b \in \mathbb{R}$ and $b > 0$.) Using de Moivre's theorem find which of the below (there may be more than one) are 6th roots of z_1 .

(a) $\text{cis}(\frac{\pi}{9}) = \text{cis}(20^\circ)$.


~~(b) $\sqrt{2} \text{cis}(\frac{4\pi}{9}) = \sqrt{2} \text{cis}(80^\circ)$~~ WRONG NORM

~~(c) $\text{cis}(\frac{8\pi}{9}) = \text{cis}(160^\circ)$~~ WRONG ANGLE

(d) $\text{cis}(\frac{10\pi}{9}) = \text{cis}(200^\circ)$

$$z_1 = \frac{-1 + \sqrt{1-4}}{2} = \frac{-1 + \sqrt{3}i}{2}$$

$$z_2 = \frac{-1 - \sqrt{3}i}{2}$$



$$z_1 = 1 \text{ cis } 120^\circ = 1 \text{ cis } (2\pi/3)$$

$$z_2 = 1 \text{ cis } 240^\circ = 1 \text{ cis } (4\pi/3)$$

$$z_1^{1/6} = 1^{1/6} \text{ cis } \left(\frac{1}{6} (120^\circ + j 360^\circ) \right) \quad j = 0, 1, 2, 3, 4, 5$$

$$= 1 \text{ cis } (20^\circ + j 60^\circ)$$

- THEY ARE
- $1 \text{ cis } (20^\circ) = 1 \text{ cis } \pi/9$
 - $1 \text{ cis } (80^\circ) = 1 \text{ cis } 4\pi/9$
 - $1 \text{ cis } (140^\circ) = 1 \text{ cis } 7\pi/9$
 - $1 \text{ cis } (200^\circ) = 1 \text{ cis } 10\pi/9$
 - $1 \text{ cis } (260^\circ) = 1 \text{ cis } 13\pi/9$
 - $1 \text{ cis } (320^\circ) = 1 \text{ cis } 16\pi/9$

3. [5 Marks] Let $\vec{w} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$, $\vec{v} = \begin{pmatrix} 3 \\ 4 \\ 5 \end{pmatrix}$ and $\vec{e}_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$ be vectors in \mathbb{R}^3 . Compute the following:

- $\vec{w} + \vec{v}$
- $|\vec{w}|, |\vec{v}|$
- $\vec{w} \cdot \vec{v}, \vec{v} \cdot \vec{e}_3$
- The angle (in degrees or radians) between \vec{v} and \vec{e}_3 . (Useful fact: $\sqrt{50} = \sqrt{2}\sqrt{25}$.)
- $\text{proj}_{\vec{v}}\vec{w}$, $\text{orth}_{\vec{v}}\vec{w}$

$$\vec{w} + \vec{v} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} + \begin{pmatrix} 3 \\ 4 \\ 5 \end{pmatrix} = \begin{pmatrix} 4 \\ 4 \\ 6 \end{pmatrix}$$

$$|\vec{w}| = \sqrt{1^2 + 0^2 + 1^2} = \sqrt{2}$$

$$|\vec{v}| = \sqrt{3^2 + 4^2 + 5^2} = \sqrt{50} = 5\sqrt{2}$$

$$\vec{w} \cdot \vec{v} = 3 + 0 + 5 = 8$$

$$\vec{v} \cdot \vec{e}_3 = 0 + 0 + 5 = 5$$

LET θ BE THE ANGLE BETWEEN \vec{v} AND \vec{e}_3

$$\cos \theta = \frac{\vec{v} \cdot \vec{e}_3}{|\vec{v}| |\vec{e}_3|} = \frac{5}{5\sqrt{2} \cdot 1} = \frac{1}{\sqrt{2}} \quad \text{SO } \theta = 45^\circ = \frac{\pi}{4}$$

$$\text{proj}_{\vec{v}} \vec{w} = \frac{\vec{v} \cdot \vec{w}}{|\vec{v}|^2} \vec{v} = \frac{8}{50} \begin{pmatrix} 3 \\ 4 \\ 5 \end{pmatrix} = \begin{pmatrix} 12/25 \\ 16/25 \\ 4/5 \end{pmatrix}$$

$$\text{ORTH}_{\vec{v}} \vec{w} = \vec{w} - \text{proj}_{\vec{v}} \vec{w} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} - \begin{pmatrix} 12/25 \\ 16/25 \\ 4/5 \end{pmatrix} = \begin{pmatrix} 13/25 \\ -16/25 \\ 1/5 \end{pmatrix}$$

4. [5 Marks] Let $\vec{u} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}$ and $\vec{v} = \begin{pmatrix} d \\ e \\ f \end{pmatrix}$ be vectors in \mathbb{R}^3 .

$$\begin{pmatrix} d \\ e \\ f \end{pmatrix} \begin{pmatrix} a \\ b \\ c \end{pmatrix}$$

- Expand the expression

$$\vec{u} \cdot (\vec{v} \times \vec{u})$$

and simplify it as much as possible (it should simplify a lot.)

- What did you just prove? (Multiple choice, one answer.)

(a) \vec{u} is orthogonal to $\vec{u} \times \vec{v}$.

(b) \vec{u} is parallel to $\vec{u} \times \vec{v}$.

(c) $\vec{u} \times \vec{v}$ always equals $\vec{0}$.

$$\begin{aligned} \begin{pmatrix} a \\ b \\ c \end{pmatrix} \cdot \begin{pmatrix} ec - bf \\ fa - dc \\ db - ea \end{pmatrix} &= aec - abf + bfa - bdc \\ &\quad + cdb - cea \\ &= ace - abf + abf - bcd \\ &\quad + bcd - ace \\ &= ace - ace + \\ &\quad abf - abf + \\ &\quad bcd - bcd \\ &= 0 \end{aligned}$$

Page intentionally left blank as extra work space