

Solutions

School of Mathematics and Statistics
Carleton University
Math. 2004A, Fall 2013
TEST 1

Any non-programmable calculator permitted, 1 blank sheet permitted for roughs

Print Name :

Student Number:

Tutorial Section (A1, A4, ...):

PART I: Multiple Choice Questions

(Choose and CIRCLE only ONE answer - No part marks here.)

- [2 marks] What value of x will make the two vectors $(x, 3)$ and $(2, 12)$ orthogonal?
(a) $x = 12$, (b) $x = -18$, (c) $x = 0$, (d) $x = -4$.
- [2 marks] Find an equation of the line through the points $(1, 2, 4)$ and $(3, -1, 6)$.
(a) $x = 1 + 2t, y = 2 - 3t, z = 4 + 2t$, (b) $x = 1 + 2t, y = 1 - t, z = -1 + 2t$, (c) $x = 1 + t, y = 1 - 2t, z = -1 + 2t$,
(d) $x = 1, y = 1 - t, z = -1 + 2t$.
- [2 marks] Find a normal vector to the plane through the points $(1, -1, 0)$, $(2, 1, 1)$ and $(-1, 0, 1)$.
(a) $(0, 1, 2)$, (b) $(3, 1, -4)$, (c) $(2, 0, -1)$, (d) $(1, -3, 5)$.
- [2 marks] Two vectors in three dimensional space are parallel if and only if their cross product is zero.
(a) TRUE, (b) FALSE, (c) It depends.
- [2 marks] Convert the equation of the circle $(x - 1)^2 + y^2 = 1$ to polar coordinates.
(a) $r = 2$, (b) $r = 3 \sin \theta$, (c) $r = 2 \cos \theta$, (d) $r = -\cos \theta$.

PART II: Show all work here and give details.

No additional pages will be accepted

- [10 marks] Find the arc length of the curve $r = 2 \sin \theta$ between the rays $\theta = 0$ and $\theta = \pi/4$.

$$\begin{aligned} \text{Arc length} &= \int_0^{\pi/4} \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta \quad \leftarrow \textcircled{5} \\ &= \int_0^{\pi/4} \sqrt{4 \sin^2 \theta + 4 \cos^2 \theta} d\theta \quad \leftarrow \textcircled{1} \\ &= 2 \int_0^{\pi/4} \sqrt{\sin^2 \theta + \cos^2 \theta} d\theta \quad \leftarrow \textcircled{1} \\ &= 2 \int_0^{\pi/4} d\theta \\ &= \pi/2 \quad \leftarrow \textcircled{2} \end{aligned}$$

7. [10 marks] Find the area under the curve $r = \sqrt{\cos \theta}$ between the rays $\theta = 0$ and $\theta = \pi/2$.

$$\begin{aligned}
 \text{Area} &= \int_0^{\pi/2} \frac{1}{2} r^2 d\theta \quad \left(= \frac{1}{2} \int_0^{\pi/2} (\cos \theta)^2 d\theta \right) \\
 &= \frac{1}{2} \int_0^{\pi/2} \cos \theta d\theta \quad \leftarrow \textcircled{2} \\
 &= \frac{1}{2} \sin \theta \Big|_0^{\pi/2} \\
 &= \left(\frac{1}{2} \right) \quad \leftarrow \textcircled{3}
 \end{aligned}$$