

Solution

For rough work, you may use back pages or else, ask me for scrap paper.

Last Name _____ First Name _____ Student Number _____

By signing below, you declare that this work was your own and that you have not copied from any other individual or other source.

Signature _____ (should match the signature that you sign on the attendance sheets.)

Questions 1–4 are multiple-choice (5 points each). You must enter the letter corresponding to each correct answer in the table below. No partial marks will be given for other work.

Problem	1 (5 pts.)	2(5 pts.)	3(5 pts.)	4(5 pt.)
Your answer	D	D	A	A
Your marks				

Question 1. Determine the derivative of $f(x) = \sin^{-1}(2x) = \arcsin(2x)$ with respect to x .

- (a) $\frac{1}{\sqrt{1-x^2}}$ (b) $\frac{2}{\sqrt{1-x^2}}$ (c) $\frac{1}{\sqrt{1-4x^2}}$ (d) $\frac{2}{\sqrt{1-4x^2}}$ (e) $\frac{9}{1-9x^2}$ (f) $\frac{9}{\sqrt{1-9x^2}}$

Using the chain rule, $f'(x) = \frac{1}{\sqrt{1-(2x)^2}} \times 2$.

Question 2. Find the slope of the tangent line to the curve $y = e^{1+x^2} \ln(1+x^2)$ at the point $x = 2$.

- (a) $\ln(2)$ (b) 1 (c) π (d) 0 (e) $e^2 \ln(2)$ (f) 1

$$y' = 2xe^{1+x^2} \ln(1+x^2) + e^{1+x^2} \frac{2x}{1+x^2}$$

$$y'(0) = 2(0)e^{1+0^2} \ln(1+0^2) + e^{1+0^2} \frac{2(0)}{1+0^2} = 0 + 0 = 0$$

Question 3. A particle moves with acceleration function $a(t) = \sin t - 3 \cos t$. Its initial velocity is $v(0) = 2$ m/s and its initial displacement is $s(0) = 5$ m. Find its position after π seconds.

- (a) $3\pi - 1$ (b) $3\pi + 1$ (c) 0 (d) π (e) $\pi + 1$ (f) $\pi - 1$

$$v(t) = -\cos t - 3 \sin t + C$$

$$2 = v(0) = -\cos 0 - 3 \sin 0 + C, \text{ so } C = 3$$

$$s(t) = -\sin t + 3 \cos t + 3t + D$$

$$5 = s(0) = -\sin 0 + 3 \cos 0 + 3(0) + D, \text{ so } D = 2$$

$$s(t) = -\sin t + 3 \cos t + 3t + 2, \text{ so } s(\pi) = -\sin \pi + 3 \cos \pi + 3\pi + 2 = 3\pi - 1.$$

Question 4. Evaluate the area between the x -axis and $f(x) = -x^2 + 4x - 3$ from $x = 1$ to $x = 3$.

- (a) $4/3$ (b) $3/4$ (c) 0 (d) $5/4$ (e) $4/5$ (f) $1/2$

$$\text{Area} = \int_1^3 -x^2 + 4x - 3 \, dx = \left[-\frac{1}{3}x^3 + 2x^2 - 3x \right]_1^3 = (-9 + 18 - 9) - \left(-\frac{1}{3} + 2 - 3 \right) = 0 - \left(-\frac{4}{3} \right) = \frac{4}{3}.$$

Questions 5-12 are long-answer (10 points each). You must clearly show all relevant steps and justify your solution to receive full marks. Clearly indicate the final answer.

Question 5. Find the derivative of the function

$$y = (\sin x)^{x^2}.$$

$$\begin{aligned} \ln y &= \ln((\sin x)^{x^2}) = x^2 \ln(\sin x) \\ \frac{y'}{y} &= 2x \ln(\sin x) + x^2 \frac{\cos x}{\sin x} \\ y' &= y(2x \ln(\sin x) + x^2 \cot x) \\ y' &= (\sin x)^{x^2} (2x \ln(\sin x) + x^2 \cot x) \end{aligned}$$

Question 6. (a) Find the linearization of $f(x) = \sqrt[3]{1+3x}$ at $a = 0$, and state the corresponding linear approximation.

$f(x) = (1+3x)^{1/3} \Rightarrow f'(x) = (1+3x)^{-2/3}$, so the linearization of f at $a = 0$ is
 $L(x) = f(0) + f'(0)(x-0) = 1^{1/3} + 1^{-2/3}x = 1 + x$. Thus, $\sqrt[3]{1+3x} \approx 1 + x$.

(b) Use part (a) to give an approximate value for $\sqrt[3]{1.03}$.

$$\sqrt[3]{1.03} = \sqrt[3]{1+3(0.01)} \approx 1 + 0.01 = 1.01.$$

Question 7. Evaluate the following limit by first recognizing the sum as a Riemann sum for a function defined on $[0, 1]$.

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \left(\sqrt[3]{\frac{1}{n}} + \sqrt[3]{\frac{2}{n}} + \sqrt[3]{\frac{3}{n}} + \cdots + \sqrt[3]{\frac{n-1}{n}} + \sqrt[3]{\frac{n}{n}} \right) \\ \lim_{n \rightarrow \infty} \frac{1}{n} \left(\sqrt[3]{\frac{1}{n}} + \sqrt[3]{\frac{2}{n}} + \sqrt[3]{\frac{3}{n}} + \cdots + \sqrt[3]{\frac{n-1}{n}} + \sqrt[3]{\frac{n}{n}} \right) = \lim_{n \rightarrow \infty} \frac{1-0}{n} \sum_{i=1}^n \sqrt[3]{\frac{i}{n}} = \int_0^1 \sqrt[3]{x} = \left[\frac{3x^{4/3}}{4} \right]_0^1 = 3/4. \end{aligned}$$

Question 8. Compute the derivative of the function given by

$$g(x) = \int_1^{x^3} \sin(1-t^2) dt.$$

By the fundamental theorem of calculus we have

$$g'(x) = (x^3)' \sin(1-(x^3)^2) - (1)' \sin(1-(1)^2) = 3x^2 \sin(1-x^6).$$

Question 9. Find the area of the region bounded by the graphs of $f(x) = x^2 + 2$, $g(x) = -x$, $x = 0$, and $x = 1$.

$$\text{Area} = \int_0^1 f(x) - g(x) dx = \int_0^1 x^2 + 2 - (-x) dx = \left[\frac{1}{3}x^3 + 2x + \frac{1}{2}x^2 \right]_0^1 = \frac{1}{3} + 2 + \frac{1}{2} = \frac{2+12+3}{6} = \frac{17}{6}.$$

Question 10. Evaluate

$$\int \frac{-4x}{(1-2x^2)^2} dx.$$

Using substitution, $u = 1 - 2x^2$, we have $du = -4x dx$ and

$$\int \frac{-4x}{(1-2x^2)^2} dx = \int \frac{1}{u^2} du = \int u^{-2} du = \frac{1}{-2+1} u^{-2+1} + C = -\frac{1}{1-2x^2} + C.$$

Question 11. Evaluate

$$\int_1^2 \frac{1}{x(1+(\ln x)^2)} dx$$

We use an integration by substitution. Substitute

$$u = \ln(x) \Rightarrow \frac{du}{dx} = \frac{1}{x} \Rightarrow dx = x du.$$

Also, we have

$$\frac{x}{u} \left| \begin{array}{c|c} 1 & 2 \\ \ln(1) = 0 & \ln(2) \end{array} \right.$$

Therefore, we get

$$\int_1^2 \frac{1}{x(1 + (\ln x)^2)} dx = \int_0^{\ln(2)} \frac{1}{1 + u^2} du = [\arctan(u)]_0^{\ln(2)} = \arctan(\ln(2)).$$

Question 12. Evaluate

$$\int_0^{\pi/2} \sin x \cos(\cos x) dx$$

Let $u = \cos x$, so $du = -\sin x dx$. When $x = 0$, $u = 1$; when $x = \pi/2$, $u = 0$. Thus,
 $\int_0^{\pi/2} \sin x \cos(\cos x) dx = \int_1^0 (-\cos u) du = [-\sin u]_1^0 = -\sin 0 - (-\sin 1) = \sin 1.$

Problem	5(10 pts.)	6(10 pts.)	7(10 pts.)	8(10 pts.)	9(10 pts.)	10(10 pts.)	11(10 pts.)	12(10 pts.)
Your marks								

Student Last,First Name: _____ Total marks: _____ out of 100

Solution

For rough work, you may use back pages or else, ask me for scrap paper.

Last Name _____ First Name _____ Student Number _____

By signing below, you declare that this work was your own and that you have not copied from any other individual or other source.

Signature _____ (should match the signature that you sign on the attendance sheets.)

Questions 1-4 are multiple-choice (5 points each). You must enter the letter corresponding to each correct answer in the table below. No partial marks will be given for other work.

Problem	1 (5 pts.)	2(5 pts.)	3(5 pts.)	4(5 pt.)
Your answer	F	C	F	B
Your marks				

Question 1. Determine the derivative of $f(x) = \sin^{-1}(3x) = \arcsin(3x)$ with respect to x
 (a) $\frac{1}{\sqrt{1-x^2}}$ (b) $\frac{2}{\sqrt{1-x^2}}$ (c) $\frac{1}{\sqrt{1-4x^2}}$ (d) $\frac{2}{\sqrt{1-4x^2}}$ (e) $\frac{9}{1-9x^2}$ (f) $\frac{3}{\sqrt{1-9x^2}}$

Using the chain rule, $f'(x) = \frac{1}{\sqrt{1-(3x)^2}} \times 3$.

Question 2. Find the slope of the tangent line to the curve $y = e^{1+x^4} \ln(1+x^4)$ at the point $x = 0$.
 (a) $\ln(2)$ (b) 1 (c) 0 (d) 0π (e) $e^2 \ln(2)$ (f) 1

$$y' = 4x^3 e^{1+x^4} \ln(1+x^4) + e^{1+x^4} \frac{4x^3}{1+x^4}$$

$$y'(0) = 4(0^3)e^{1+0^4} \ln(1+0^4) + e^{1+0^4} \frac{4(0^3)}{1+0^4} = 0 + 0 = 0$$

Question 3. A particle moves with acceleration function $a(t) = \cos t - 3 \sin t$. Its initial velocity is $v(0) = 2$ m/s and its initial displacement is $s(0) = 5$ m. Find its position after π seconds.

(a) $3\pi - 1$ (b) $3\pi + 1$ (c) $\pi - 7$ (d) π (e) $\pi + 7$ (f) $7 - \pi$

$$v(t) = \sin t + 3 \cos t + C$$

$$2 = v(0) = \sin 0 + 3 \cos 0 + C, \text{ so } C = -1$$

$$s(t) = -\cos t + 3 \sin t - t + D$$

$$5 = s(0) = -\cos 0 + 3 \sin 0 - 0 + D, \text{ so } D = 6$$

$$s(t) = -\cos t + 3 \sin t - t + 6, \text{ so } s(\pi) = -\cos \pi + 3 \sin \pi - \pi + 6 = 1 + 0 - \pi + 6 = 7 - \pi.$$

Question 4. Evaluate the area between the x -axis and $f(x) = -x^2 + 4x - 3$ from $x = 1$ to $x = 3$.
 (a) $3/4$ (b) $4/3$ (c) $5/4$ (d) 0 (e) $4/5$ (f) $1/2$

$$\text{Area} = \int_1^3 -x^2 + 4x - 3 dx = \left[-\frac{1}{3}x^3 + 2x^2 - 3x \right]_1^3 = (-9 + 18 - 9) - \left(-\frac{1}{3} + 2 - 3 \right) = 0 - \left(-\frac{4}{3} \right) = \frac{4}{3}.$$

Questions 5-12 are long-answer (10 points each). You must clearly show all relevant steps and justify your solution to receive full marks. Clearly indicate the final answer.

Question 5. Find the derivative of the function

$$y = (\cos x)^{x^3}.$$

$$\ln y = \ln((\cos x)^{x^3}) = x^3 \ln(\cos x)$$

$$\frac{y'}{y} = 3x^2 \ln(\cos x) + x^3 \frac{-\sin x}{\cos x}$$

$$y' = y(3x^2 \ln(\cos x) - x^3 \tan x)$$

$$y' = (\cos x)^{x^3} (3x^2 \ln(\cos x) - x^3 \tan x)$$

Question 6. (a) Find the linearization of $f(x) = \sqrt[4]{1+4x}$ at $a = 0$, and state the corresponding linear approximation.

$$f(x) = (1+4x)^{1/4} \Rightarrow f'(x) = (1+4x)^{-3/4}, \text{ so the linearization of } f \text{ at } a = 0 \text{ is}$$

$$L(x) = f(0) + f'(0)(x-0) = 1^{1/4} + 1^{-3/4}x = 1 + x. \text{ Thus, } \sqrt[4]{1+4x} \approx 1 + x.$$

(b) Use part (a) to give an approximate value for $\sqrt[4]{1.04}$.

$$\sqrt[4]{1.04} = \sqrt[4]{1+4(0.01)} \approx 1 + 0.01 = 1.01.$$

Question 7. Evaluate the following limit by first recognizing the sum as a Riemann sum for a function defined on $[0, 1]$.

$$\lim_{n \rightarrow \infty} \frac{1}{n} \left(\sqrt[5]{\frac{1}{n}} + \sqrt[5]{\frac{2}{n}} + \sqrt[5]{\frac{3}{n}} + \cdots + \sqrt[5]{\frac{n-1}{n}} + \sqrt[5]{\frac{n}{n}} \right)$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \left(\sqrt[5]{\frac{1}{n}} + \sqrt[5]{\frac{2}{n}} + \sqrt[5]{\frac{3}{n}} + \cdots + \sqrt[5]{\frac{n-1}{n}} + \sqrt[5]{\frac{n}{n}} \right) = \lim_{n \rightarrow \infty} \frac{1-0}{n} \sum_{i=1}^n \sqrt[5]{\frac{i}{n}} = \int_0^1 \sqrt[5]{x} = \left[\frac{5x^{6/5}}{6} \right]_0^1 = 5/6.$$

Question 8. Compute the derivative of the function given by

$$g(x) = \int_0^{x^2} \cos(1-t^2) dt.$$

By the fundamental theorem of calculus we have

$$g'(x) = (x^2)' \cos(1-x^2) - (0)' \cos(1-t^2) = 2x \cos(1-x^2).$$

Question 9. Find the area of the region bounded by the graphs of $f(x) = x^3 + 2$, $g(x) = -x$, $x = 0$, and $x = 1$.

$$\text{Area} = \int_0^1 f(x) - g(x) dx = \int_0^1 x^3 + 2 - (-x) dx = \left[\frac{1}{4}x^4 + 2x + \frac{1}{2}x^2 \right]_0^1 = \frac{1}{4} + 2 + \frac{1}{2} = \frac{11}{4}.$$

Question 10. Evaluate

$$\int \frac{-4x}{(1-2x^2)^3} dx.$$

Using substitution, $u = 1 - 2x^2$, we have $du = -4x dx$ and

$$\int \frac{-4x}{(1-2x^2)^3} dx = \int \frac{1}{u^3} du = \int u^{-3} du = \frac{1}{-3+1} u^{-3+1} + C = \frac{-1}{2(1-2x^2)^2} + C.$$

Question 11. Evaluate

$$\int \frac{1}{e^x(1+(e^x)^2)} dx.$$

We use an integration by substitution. Substitute

$$u = e^x \Rightarrow \frac{du}{dx} = e^x \Rightarrow dx = \frac{du}{e^x}.$$

Also, we have

$$\frac{x}{u} \Big|_{e^0=1}^1 \Big|_e.$$

Therefore, we get

$$\int_0^1 \frac{e^x}{(1+(e^x)^2)} dx = \int_1^e \frac{1}{1+u^2} du = [\arctan(u)]_1^e = \arctan(e) - \arctan(1) = \arctan(e) - \frac{\pi}{4}.$$

Question 12. Evaluate

$$\int_0^{\pi/2} \cos x \sin(\sin x) dx$$

Let $u = \sin x$, so $du = \cos x dx$. When $x = 0$, $u = 0$; when $x = \pi/2$, $u = 1$. Thus,
 $\int_0^{\pi/2} \cos x \sin(\sin x) dx = \int_0^1 \sin u du = [-\cos u]_0^1 = -\cos 1 - (-\cos 0) = 1 - \cos 1.$

Problem	5(10 pts.)	6(10 pts.)	7(10 pts.)	8(10 pts.)	9(10 pts.)	10(10 pts.)	11(10 pts.)	12(10 pts.)
Your marks								

Student Last,First Name: _____ Total marks: _____ out of 100