

CARLETON UNIVERSITY

FINAL EXAMINATION
MATH 1004 A, B, C, D, E, F
 December 2013

DURATION: 3 HOURS

Department Name and Course Number: School of Mathematics and Statistics,
 MATH 1004 A, B, C, D, E, F.

Course Instructor(s): Dr. A.B. Mingarelli (Sect. A), Dr. Y. Gao (Sect. B), Mr. M. Blenkinsop (Sect. C, D), Dr. B. Brimacombe (Sect. E), Dr. Z. Montazeri (Sect. F).

AUTHORIZED MEMORANDA

NON-PROGRAMMABLE CALCULATOR PERMITTED.

BLANK SHEETS OF PAPER PROVIDED BY THE UNIVERSITY

This exam may be released to the Library and may be taken away by the student.

1. Please verify that you are in possession of a Scantron FORM
2. Please **fill in your COURSE CODE** (e.g., MATH 1004) and **COURSE SECTION** (e.g., A, B, C, D, E, F), **YOUR NAME** and **YOUR STUDENT NUMBER** where required on the Scantron form.
3. **The examination consists of two sheets of legal size paper.** It is out of a total of 100 and consists of 25 multiple choice questions each worth 4 marks **Please fill in only one answer on your Scantron sheets with a pencil** as there is only one answer to any given question. Circling two or more answers to any question invalidates that question (*i.e.*, you get 0 marks for that question).

Return only the duly completed Scantron form, not the examination nor your work.

1. [4 marks] Let $f(x) = |x - 1| + |x - 3|$. Calculate $L = \lim_{h \rightarrow 0^+} \frac{f(1+h) - f(3)}{h}$.
 (a) $L = 0$ (b) $L = 1$ (c) $L = -1$ (d) This limit does not exist
2. [4 marks] Let $f(x) = \frac{1 - \cos x}{x^2}$, for $x \neq 0$, and $f(x) = A$, for $x = 0$. What value of A will make f continuous at $x = 0$?
 (a) $A = 0$ (b) $A = 1/2$ (c) $A = -1$ (d) $A = 1$.
3. [4 marks] Evaluate $L = \lim_{x \rightarrow 3} \frac{x^2 - 2x - 3}{x^2 - 9}$.
 (a) $L = 0$ (b) $L = \frac{3}{2}$ (c) $L = \frac{2}{3}$ (d) This limit does not exist
4. [4 marks] Let $f(x) = \frac{\sin(3x)}{\sin(2x)}$. Evaluate $L = \lim_{x \rightarrow 0} f(x)$.
 (a) $L = 0$ (b) $L = \frac{3}{2}$ (c) $L = \frac{2}{3}$ (d) This limit does not exist
5. [4 marks] Two functions f, g are defined by $f(x) = 3x^2$ and $g(x) = \cos x$. What is the value of their composition $f(g(0))$?
 (a) -3 (b) 3 (c) -3.2 (d) 0

6. [4 marks] Find the derivative of the function $y = \frac{8x}{\ln(5x+1)}$.
- (a) $\frac{8(5x+1)\ln(5x+1) - 40x}{(5x+1)(\ln(5x+1))^2}$ (b) $\frac{1}{\ln(40x)}$ (c) $\frac{8\ln(5x+1) - 40x}{\ln(5x+1)(5x+1)^2}$ (d) $\frac{8}{5\ln(5x+1)}$
7. [4 marks] Find the derivative of the function $y = 5x^2e^{3x}$.
- (a) $10xe^{3x}(2x+3)$ (b) $5xe^{3x}(2x+3)$ (c) $10ex^{3x}(3x+2)$ (d) $5xe^{3x}(3x+2)$
8. [4 marks] Find the derivative of the function $y = \ln(e^{x^2} + 1)$.
- (a) $\frac{2xe^{x^2}}{e^{x^2} + 1}$ (b) $\frac{2x}{e^{x^2}}$ (c) $\frac{2xe^{x^2}}{\ln(e^{x^2} + 1)}$ (d) $\frac{2e^{x^2}}{(e^{x^2} + 1)^2}$
9. [4 marks] Find any local maximum or minimum points of the given function. $y = x^3 - 3x^2 + 1$.
- (a) Minimum at $(0, 1)$, maximum at $(2, -3)$ (b) Maximum at $(0, 1)$, minimum at $(2, -3)$ (c) Maxima at $(-2, -19)$ and $(0, 1)$, minimum at $(2, -3)$ (d) Minimum at $(2, -3)$
10. [4 marks] Which of the following statements is true?
- (a) $f(x) = 2e^x$ is concave down for all x , and has no points of inflection.
 (b) $f(x) = x^5 + 1$ is concave up for all x , and has no points of inflection.
 (c) $f(x) = x^2 + 5$ is concave up for $x < 0$, concave down for $x > 0$, and has a point of inflection at $(0, 5)$.
 (d) $f(x) = (x - 5)^3$ is concave down for $x < 5$, concave up for $x > 5$, and has a point of inflection at $(5, 0)$.
11. [4 marks] Evaluate $\int \frac{\sec^2(\ln x)}{x} dx$.
- (a) $\tan(\ln x) + C$ (b) $\ln(\sec x) + C$, (c) $2\sec(\ln x) + C$ (d) $\ln(\tan x) + C$
12. [4 marks] Evaluate the definite integral $\int_0^\pi \sin^2\left(\frac{x}{2}\right) \cos^2\left(\frac{x}{2}\right) dx$
- (a) $\frac{\pi}{6}$ (b) $\frac{\pi}{2}$ (c) $\frac{\pi}{4}$ (d) $\frac{\pi}{8}$
13. [4 marks] Evaluate $I = \int e^{4x} \cos\left(\frac{x}{2}\right) dx$.
- (a) $\frac{1}{12}e^{4x} \left(3 \sin\left(\frac{x}{2}\right) + 14 \cos\left(\frac{x}{2}\right)\right) + C$ (b) $\frac{1}{23}e^{4x} \left(2 \sin\left(\frac{x}{2}\right) + 3 \cos\left(\frac{x}{2}\right)\right) + C$
 (c) $\frac{1}{65}e^{4x} \left(2 \sin\left(\frac{x}{2}\right) + 16 \cos\left(\frac{x}{2}\right)\right) + C$ (d) $\frac{1}{5}e^{4x} \left(\sin\left(\frac{x}{2}\right) - \cos\left(\frac{x}{2}\right)\right) + C$
14. [4 marks] Evaluate the definite integral $\int_0^3 e^{x/3}(x^2 + 2x) dx$.
- (a) 0 (b) $27e - 36$ (c) $e - 1$ (d) $21e + 40$
15. [4 marks] Evaluate the definite integral $\int_1^e (x \ln x)^2 dx$.
- (a) $\frac{e}{4}$ (b) $\frac{e^3 - 1}{2}$ (c) $\frac{5e^3 - 2}{27}$ (d) $\frac{e^2 + 1}{6}$
16. [4 marks] Evaluate $I = \int \frac{4}{x^4 - 1} dx$.
- (a) $\ln|x - 1| - \ln|x + 1| - 2 \tan^{-1}(x) + C$ (b) $\ln|x^2 + 1| + 2 \tan^{-1}(x) + C$
 (c) $\ln|x - 1| - 4 \ln|x + 1| - 2 \tan^{-1}(x) + C$ (d) $2 \ln|x - 1| + \ln|x + 1| + \tan^{-1}(x) + C$
17. [4 marks] Let $f(x) = \sin(\sin 3x)$. Evaluate $f'(\pi/2)$. In other words, find the derivative of f at $x = \pi/2$.
- (a) $f'(\pi/2) = 0$ (b) $f'(\pi/2) = 1$ (c) $f'(\pi/2) = 2$ (d) $f'(\pi/2) = 3$

18. [4 marks] Evaluate the following limit: $L = \lim_{x \rightarrow 0} \frac{\arcsin(5x)}{x^2}$
 (a) $L = 5$ (b) $L = \frac{1}{5}$ (c) $L = 0$ (d) This limit does not exist
19. [4 marks]. Given that f is such that its inverse F exists, $f'(-5) = 4$, $F(2) = -5$, find the value of the derivative of F at $x = 2$.
 (a) 4 (b) $1/4$ (c) 5 (d) $1/5$
20. [4 marks] Let y be given implicitly as a differentiable function of x by $2x = xy + y^2$. Then the slope of the tangent line to the curve $y = y(x)$ at the point (x, y) where $x = 1, y = 1$ is equal to:
 (a) 2 (b) $1/2$ (c) 3 (d) $1/3$
21. [4 marks] Let $f(x) = 2|x - 5|$. Calculate $L = \lim_{h \rightarrow 0} \frac{f(5+h) - f(5)}{h}$.
 (a) $L = 0$ (b) $L = 5$ (c) $L = -5$ (d) This limit does not exist.
22. [4 marks] Let $f(x) = \sqrt{x^2 + 4}$. Evaluate $f''(0)$. In other words, find the second derivative of f at $x = 0$.
 (a) $f''(0) = 4$ (b) $f''(0) = 0$ (c) $f''(0) = 1/2$ (d) $f''(0)$ does not exist
23. [4 marks] Find an expression for the volume of the solid of revolution obtained by rotating the region in the first quadrant bounded by the curve defined by $y = \cos x$ between $x = 0$ and $x = \pi/2$ about the y -axis.
 (a) $\pi \int_0^{\pi/2} x^2 \cos x \, dx$ (b) $\int_0^{\pi/2} \cos x \, dx$ (c) $\int_0^{\pi/2} x \sin x \, dx$ (d) $2\pi \int_0^{\pi/2} x \cos x \, dx$
24. [4 marks] Evaluate the improper integral $\int_0^\infty 3x^2 e^{-x} \, dx$.
 (a) 12 (b) 2 (c) 6 (d) 1
25. [4 marks] Find the area of the region bounded by the curves $y = x^2 + 1$ and $y = 5$.
 (a) 16 (b) $\frac{8}{3}$ (c) $\frac{32}{3}$ (d) $\frac{3}{2}$

[Total: 100 marks]

END OF THE EXAMINATION.