



Université d'Ottawa · University of Ottawa

Faculté des science
Chimie

Faculty of Science
Chemistry

CHM 1301

DATE: December 9, 2013
9:30 am

Dr. Rashmi Venkateswaran

TIME: 3 hours

Page 1 of 19

FULL STUDENT NAME: (Print) _____

STUDENT NUMBER: _____

INSTRUCTIONS:

1. Write your answers on the exam. Please **BOLD** or **HIGHLIGHT** your final answer.
2. You are allowed to use electronic calculators during the exam (TI 30X, TI 34, TI 36, Casio 290 or 300).
3. Equations and constants are given on pages 17 and 18. A periodic table of the elements is included (page 19).
4. **Use the correct number of significant figures at all times.**
5. Marks allotted to each question are shown in brackets. The exam is out of 100.
7. Use the back of sheets for rough work. Three extra (blank) pages (14, 15 and 16) have also been included.
8. You are advised to read through the exam paper before you begin to write. Answer first those questions that you can do right away, proceeding to more difficult questions thereafter. Allow about ten minutes near the end of the exam period to read quickly through your answers, checking for obvious errors. **Good luck!**

Quest.	1	2	3	4	5	6	7	8	9	10	
Marks	/10	/10	/10	/10	/10	/10	/12	/8	/10	/10	/100

- (10) 1. A microwave oven operating at 1.22×10^8 nm is used to heat 150.0 mL of water (roughly the volume of a tea cup) from 20.0 °C to 95.0 °C. Calculate the number of photons needed if 92.0 percent of microwave energy is converted to the thermal energy of water. ($d_{\text{water}} = 1.00$ g/mL; $c_{\text{water}} = 4.184$ J/g°C)

[5]

(b) In a photoelectric effect experiment, photons having an energy of 361.3 kJ/mol are absorbed by a metal and the maximum kinetic energy of the resulting electrons is $KE = 162.6$ kJ/mol.

i) Calculate the binding energy (work function) of ONE electron in the metal.

ii) Calculate the wavelength of the light.

[5]

iii) Calculate the velocity of the electrons.

- (10) 2. (a) The carbon dioxide exhaled by astronauts must be removed from the spacecraft atmosphere. One way to do this is to react the gaseous carbon dioxide with solid lithium hydroxide, which results in the formation of solid lithium carbonate and liquid water. The carbon dioxide output of an astronaut is about 1.0 kg/day. What is the minimum mass of lithium hydroxide required for a six-day space shuttle flight involving five astronauts?

[5]

- (b) The combustion of what volume of ethane (C_2H_6) measured at $23.0\text{ }^\circ\text{C}$ and 752 mm Hg would be required to heat 855 g of water from $25.0\text{ }^\circ\text{C}$ to $98.0\text{ }^\circ\text{C}$? $c_{\text{water}} = 4.184\text{ J/g}^\circ\text{C}$; $\Delta_f H^\circ (CO_2) = -393.5\text{ kJ/mol}$; $\Delta_f H^\circ (H_2O(l)) = -285.8\text{ kJ/mol}$; $\Delta_f H^\circ (C_2H_6) = -84.7\text{ kJ/mol}$

[5]

(10) 3. Iodine forms three compounds with chlorine: ICl , ICl_3 , and ICl_5 .

(a) Showing all your work, determine the molecular geometries of all three molecules.

[6]

(b) What are the shapes of all three molecules? Draw them.

[3]

(c) What are the main angles in each of the three molecules? Are they polar?

[1]


- (10) 4. (a) When heated, lithium reacts with nitrogen to form lithium nitride:



What is the theoretical yield of Li_3N when 12.3 g of Li is heated with 33.6 g of N_2 ? If the actual yield of Li_3N is 5.89 g, what is the percent yield of the reaction?

[5]


- (b) A sample of zinc metal reacts completely with an excess of hydrochloric acid:



The hydrogen gas produced is collected over water at 25.0 °C. The volume of the gas is 7.80 L and the pressure is 0.980 atm. Calculate the mass of zinc metal consumed in the reaction.

[5]

- (10) 5. (a) The following initial rate information was collected at 25°C for the reaction:



Experiment	$[\text{I}^-]_0$ (mol/L)	$[\text{S}_2\text{O}_8^{2-}]_0$ (mol/L)	Initial Rate (mol/L·s)
1	0.125	0.150	4.4×10^{-2}
2	0.375	0.150	1.3×10^{-1}
3	0.125	0.050	1.5×10^{-2}

- i) Determine the rate law expressed by this data.

[4]

- ii) Calculate the rate constant with the CORRECT units.



- b) The light-emitting decay of excited mercury atoms is first order, with a half-life of 4.20×10^{-7} s. A sample contains 4.5×10^{-6} mol/L of excited mercury atoms.

- i) At what time will the concentration be 4.5×10^{-7} mol/L?

[2]

- ii) What will the concentration of mercury atoms be after 2.5×10^{-6} s?

b) The reaction of CO with Cl_2 gives phosgene (COCl_2), a nerve gas used in World War I. Even though the stoichiometry is simple, the mechanism has several steps:



- i) Show that the elementary reactions of the proposed mechanism can be added to provide a balanced equation for the reaction.

[1]

- ii) Identify any intermediates in the mechanism.

[1]

- iii) What is the rate law yielded by this mechanism?

[2]

- (10) 6. (a) The solubility of calcium phosphate in water is 3.5×10^{-5} g/L. Use this information to calculate K_{sp} for the salt.

[5]

- (b) What mass of BaF_2 ($K_{sp} = 1.8 \times 10^{-7}$) will dissolve in 0.500 L of 0.100 mol/L NaF solution?

[5]

(12) 7. (a) Consider the titration of 30.0 mL of 0.200 mol/L HCOONa (aq) with 6.0 mol/L HCl (aq). (K_a for HCOOH is 1.8×10^{-4})

i) What is the original pH of the solution of HCOONa?

ii) What is the pH of the solution halfway through the titration?

[6]


iii) What is the pH at the equivalence point?

- (b) What is the pH of a buffer solution containing 35.0 g of NH_4Cl dissolved in 1.00 L of 1.00 mol/L NH_3 ?

[6]

- ii) Calculate the pH of the system if 5.0 mL of 12.0 mol/L HCl is added to the buffer.

- (8) 8. Hydrogen fluoride is a highly reactive gas. It has many industrial uses, but the most familiar property of HF is its ability to react with glass. As a result, HF is used to etch glass and frost the inner surfaces of light bulbs. Hydrogen fluoride gas must be stored in stainless steel containers and aqueous solutions must be stored in plastic bottles. Hydrogen fluoride can be produced H_2 (g), and F_2 (g):



In a particular experiment, 3.00 bar of hydrogen gas and 2.00 bar of fluorine gas are added to a 1.50 L flask at 350 K. Calculate the equilibrium partial pressures of all species.

[5]

- (c) A coin dealer, offered a rare silver coin, suspected that it might be a counterfeit nickel copy. The dealer heated the coin, of mass 15.5 g, to 100.0 °C in boiling water and then dropped the hot coin into 21.5 g of water in a coffee cup calorimeter at 15.5 °C. The temperature of the water rose to 21.5 °C. Was the coin made of silver or nickel?

$$c_{\text{Ag}} = 0.240 \text{ J/g}^\circ\text{C}$$

$$c_{\text{Ni}} = 0.440 \text{ J/g}^\circ\text{C}$$

[3]

- (10) 9. Police officers confiscate a packet of white powder that they believe contains heroin. Purification by a forensic chemist yields a 38.70 mg sample for combustion analysis. This sample given 97.46 mg of carbon dioxide and 20.81 mg of water. A second sample is analysed for its nitrogen content, which is found to be 3.8%. Show by calculations whether these data are consistent with the formula for heroin, $C_{21}H_{22}NO_5$.

- (10) 10. Your parents have offered to pay your tuition for the fall semester of University if you can explain to them why you have selected the courses you chose and why they will be potentially helpful in getting you a job. You, of course, choose Chemistry to demonstrate how important it is to our lives today. Using **any one** of the topics covered during the semester, write a letter of minimum length 1 page and using at least 3 chemical concepts within your chosen topic (clearly explained and not just repeated from class) to demonstrate why a knowledge of chemistry is essential in our daily lives.

Equations and Constants

Gas Laws

$$PV = nRT$$

$$P_{\text{Total}} = P_1 + P_2 + P_3 + \dots$$

$$d = m/V = P(\text{MM}) / RT$$

$$KE = (1/2)mv_{\text{av}}^2$$

$$u_{\text{rms}} = \sqrt{\frac{3RT}{M}}$$

$$\frac{\text{Rate}_A}{\text{Rate}_B} = \sqrt{\frac{M_b}{M_a}}$$

$$P + \frac{n^2 a}{V^2} (V - nb) = nRT$$

Equilibrium

$$K_p = K(RT)^{\Delta n}$$

$$K = \frac{[C]^c [D]^d}{[A]^a [B]^b}$$

$$K_{\text{sp}} = [C]^c [D]^d$$

Acid/Base

$$pOH = -\log [OH^-]$$

$$pH = -\log [H^+]$$

$$pH + pOH = 14$$

$$K_a \cdot K_b = K_w$$

$$pH = pK_a + \log \frac{[A^-]}{[HA]}$$

$$pH = \frac{pK_{a1} + pK_{a2}}{2}$$

$$C_1 V_1 = C_2 V_2$$

$$\ln \left(\frac{K_1}{K_2} \right) = \frac{\Delta H^\circ}{R} \left(\frac{1}{T_2} - \frac{1}{T_1} \right)$$

Electrochemistry

$$\Delta G^\circ = -nFE^\circ$$

$$E = E^\circ - \frac{RT}{nF} \ln(Q)$$

$$\text{Or } E = E^\circ - \frac{0.059}{n} \log(Q) \quad \text{at } 25^\circ\text{C}$$

$$Q = I \cdot t$$

Quantum Mechanics

$$\lambda \cdot \nu = c$$

$$E = h \cdot \nu$$

$$m = \frac{h}{\lambda c}$$

$$E = \frac{1}{2} m u^2$$

$$\lambda = h/mu$$

$$\text{Energy of state} = -2.178 \times 10^{-18} \text{ J} / n^2$$

$$\Delta x \cdot \Delta p \geq h / 4\pi$$

$$E = -C(1/n^2)$$

Liquids and Colligative Properties

$$\ln(P_1/P_2) = \Delta H^\circ / R (1/T_2 - 1/T_1)$$

$$P_{\text{solution}} = X_{\text{solvent}} \cdot P_{\text{solvent}}^\circ$$

$$\Delta T_{\text{BP}} = K_{\text{BP}} \cdot m$$

$$\Delta T_{\text{FP}} = K_{\text{FP}} \cdot m$$

$$\Delta T = K \cdot m \cdot i$$

$$\Pi = cRT$$

$$\text{molality} = n(\text{solute})/m(\text{solvent (kg)})$$

General INFORMATION

$$n = m/\text{MM}$$

$$d = m/V$$

$$C = n/V$$

$$1 \text{ atm} = 760 \text{ Torr} = 760 \text{ mm Hg} = 1.01325 \times 10^5 \text{ Pa} = 1.01325 \text{ bar} = 101.325 \text{ kPa}$$

Thermodynamics	Kinetics
$\Delta U = q$	Rate = $k [A]^x[B]^y[C]^z$
$w_{\text{system}} = -P\Delta V$	Rate = k
$\Delta E = q + w$	Rate = $k[A]$
$\Delta H = \Delta E + P\Delta V$	Rate = $k[A]^2$
$q_p = \Delta E + P\Delta V$	$[A] = -kt + [A]_0$
$\Delta E = nC_v\Delta T$	$\ln[A] = -kt + \ln[A]_0$
$\Delta H = q_p = m C_p \Delta T$	$1/[A] = kt + 1/[A]_0$
$C_p = C_v + R$	$t = [A]_0 / 2k$
$\Delta H_{\text{rxn}}^\circ = \Delta n_p \Delta H_f^\circ(\text{products}) - \Delta n_r \Delta H_f^\circ(\text{reactants})$	$\ln 2 = kt$
$q_{\text{rev}} = -w_{\text{max}} = nRT \ln(V_2/V_1)$	$k = Ae^{-E_a/RT}$
$\Delta S = q_{\text{rev}} / T$	$\ln\left(\frac{k_1}{k_2}\right) = \frac{E_a}{R} \left(\frac{1}{T_2} - \frac{1}{T_1}\right)$
$\Delta S_{T_1-T_2} = nC_p \ln(T_2/T_1)$	Bonding
$\Delta S_{T_1-T_2} = nC_v \ln(T_2/T_1)$	$DE = k(Q_1Q_2 / r)$
$\Delta S_{\text{surroundings}}^\circ = \frac{q_{\text{surroundings}}}{T} = \frac{-\Delta H_{\text{sys}}}{T}$	$\Delta H_{\text{rxn}} = \sum n_p D(\text{reactants}) - \sum n_r D(\text{products})$
$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$	Formal Charge = #valence e^- in free atom - #lone pair e^- - 1/2(# bonding e^-)
$\Delta G = \Delta G^\circ + RT \ln(Q)$	
$\Delta G^\circ = -RT \ln(K)$	

Data For Water and Other Constants


Density $d_{\text{water}} = 1.00 \text{ g/mL (25}^\circ\text{C)}$	$c_{\text{water}} = 4.184 \text{ J g}^{-1} \text{ }^\circ\text{C}^{-1}$ (liquid)
Avogadro's Number N	$6.022 \times 10^{23} \text{ mol}^{-1}$
Boltzmann's constant k	$1.30866 \times 10^{-23} \text{ J}\cdot\text{K}^{-1}$
Faraday's constant F	96,485 $\text{C}\cdot\text{mol}^{-1}$
Gas constant R	8.31451 $\text{J}\cdot\text{K}^{-1}\cdot\text{mol}^{-1}$
	0.08206 $\text{L}\cdot\text{atm}\cdot\text{K}^{-1}\cdot\text{mol}^{-1}$
Mass of Electron m_e	$9.10938188 \times 10^{-31} \text{ kg}$
Planck's constant h	$6.62608 \times 10^{-34} \text{ J}\cdot\text{s}$
Speed of Light c	$2.99792458 \times 10^8 \text{ m}\cdot\text{s}^{-1}$

$$K_w = 1.00 \times 10^{-14}$$

$$\text{General Information: } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Prof Mokeur's Periodic Table

																		18 VIII A	
																		He 2 4.002602 - 0 Helium	
																		17 VII A	
																		F 9 18.998403 3.98 -1 Fluorine	
																		16 VI A	
																		O 8 15.9994 3.44 -2 Oxygen	
																		15 V A	
																		N 7 14.0067 3.04 5,3 Nitrogen	
																		14 IV A	
																		C 6 12.0107 2.55 4 Carbon	
																		13 III A	
																		B 5 10.811 2.04 3 Boron	
																		Al 13 26.981538 1.61 3 Aluminum	
																		12 IIB	
																		Zn 30 65.38 1.65 2 Zinc	
																		11 IB	
																		Cu 29 63.546 1.9 2,1 Copper	
																		10 VIII	
																		Ni 28 58.6934 1.88 3,2 Nickel	
																		9 VII B	
																		Co 27 58.933195 1.88 3,2 Cobalt	
																		8 VI B	
																		Fe 26 55.845 1.83 3,2 Iron	
																		7 VIB	
																		Mn 25 54.938045 1.55 7,4,2 Manganese	
																		6 V B	
																		Cr 24 51.9961 1.66 6,3 Chromium	
																		5 VB	
																		V 23 50.9415 1.63 5 Vanadium	
																		4 IV B	
																		Ti 22 47.867 1.54 4 Titanium	
																		3 IIIB	
																		Sc 21 44.955912 1.36 3 Scandium	
																		2 IIA	
																		Be 4 9.012182 1.57 2 Beryllium	
																		1 IA	
																		Li 3 6.941 0.98 1 Lithium	
																		H 1 1.00794 2.20 1 Hydrogen	
																		Na 11 22.989769 0.93 1 Sodium	
																		Mg 12 24.3050 1.31 2 Magnesium	
																		K 19 39.0983 0.82 1 Potassium	
																		Ca 20 40.078 1.0 2 Calcium	
																		Sr 38 87.62 0.95 2 Strontium	
																		Rb 37 85.4678 0.82 1 Rubidium	
																		Cs 55 132.90545 0.79 1 Cesium	
																		Ba 56 137.327 0.89 2 Barium	
																		Ra 88 226.0254 0.7 1 Radium	
																		Fr 87 223.0197 0.7 1 Francium	
																		Ac 89 227.0278 1.1 3 Actinium	
																		Rf 104 261.11 - - - Rutherfordium	
																		Db 105 262.11 - - - Dubnium	
																		Sg 106 263.12 - - - Seaborgium	
																		Bh 107 262.12 - - - Bohrium	
																		Hs 108 264 - - - Hassium	
																		Mt 109 266.1378 - - - Meitnerium	
																		Ds 110 269 - - - Darmstadtium	
																		Rg 111 272 - - - Roentgenium	
																		Cn 112 277 - - - Copernicium	
																		Uut 113 284 - - - Ununtrium	
																		Fl 114 289 - - - Flerovium	
																		Lv 116 282 - - - Livermorium	
																		Uus 117 293 - - - Ununseptium	
																		Uuo 118 294 - - - Ununoctium	
																		Rn 86 222.0176 2.2 0 Radon	
																		At 85 208.9871 2.2 -1 Astatine	
																		Po 84 209 2.0 4,2 Polonium	
																		Bi 83 208.98040 2.02 3 Bismuth	
																		Pb 82F 207.2 2.33 4,2 Lead	
																		Tl 81 204.3833 1.62 3,1 Thallium	
																		Hg 80 200.59 2.0 2,1 Mercury	
																		Au 79 196.96657 2.54 3 Gold	
																		Pt 78 195.084 2.28 4,2 Platinum	
																		Ir 77 182.217 2.2 4,3 Iridium	
																		Os 76 190.23 2.2 4 Osmium	
																		Re 75 186.207 1.9 4 Rhenium	
																		W 74 183.84 2.36 6,4 Tungsten	
																		Ta 73 180.94788 1.5 5 Tantalum	
																		Nb 41 92.90638 1.6 5 Niobium	
																		Mo 42 95.96 2.16 6,4 Molybdenum	
																		Tc 43 98.90625 1.9 7,4 Technetium	
																		Ru 44 101.07 2.2 4,3 Ruthenium	
																		Rh 45 102.90550 2.28 3 Rhodium	
																		Pd 46 106.42 2.2 4,2 Palladium	
																		Ag 47 107.8682 1.93 1 Silver	
																		Cd 48 112.411 1.69 2 Cadmium	
																		In 49 114.818 1.78 3 Indium	
																		Sn 50 118.710 2.05 5,3 Tin	
																		Sb 51 121.760 2.1 6,4,2 Antimony	
																		Te 52 127.60 2.1 6,4,2 Tellurium	
																		I 53 126.90447 2.7 5,3,1 Iodine	
																		Xe 54 131.293 2.60 4,2 Xenon	
																		Kr 36 83.798 3.00 0,2 Krypton	
																		Br 35 79.904 2.96 5,3,1 Bromine	
																		Se 34 78.96 2.55 6,4,2 Selenium	
																		As 33 74.92160 2.18 5,3 Arsenic	
																		Ge 32 72.64 2.01 4,2 Germanium	
																		Ga 31 69.723 1.81 3 Gallium	
																		Zn 30 65.38 1.65 2 Zinc	
																		Cu 29 63.546 1.9 2,1 Copper	
																		Ni 28 58.6934 1.88 3,2 Nickel	
																		Co 27 58.933195 1.88 3,2 Cobalt	
																		Fe 26 55.845 1.83 3,2 Iron	
																		Mn 25 54.938045 1.55 7,4,2 Manganese	
																		Cr 24 51.9961 1.66 6,3 Chromium	
																		V 23 50.9415 1.63 5 Vanadium	
																		Ti 22 47.867 1.54 4 Titanium	
																		Sc 21 44.955912 1.36 3 Scandium	
																		Ca 20 40.078 1.0 2 Calcium	
																		Sr 38 87.62 0.95 2 Strontium	
																		Rb 37 85.4678 0.82 1 Rubidium	
																		Cs 55 132.90545 0.79 1 Cesium	
																		Ba 56 137.327 0.89 2 Barium	
																		Ra 88 226.0254 0.7 1 Radium	
																		Fr 87 223.0197 0.7 1 Francium	
																		Ac 89 227.0278 1.1 3 Actinium	
																		Rf 104 261.11 - - - Rutherfordium	
																		Db 105 262.11 - - - Dubnium	
																		Sg 106 263.12 - - - Seaborgium	
																		Bh 107 262.12 - - - Bohrium	
																		Hs 108 264 - - - Hassium	
																		Mt 109 266.1378 - - - Meitnerium	
																		Ds 110 269 - - - Darmstadtium	
																		Rg 111 272 - - - Roentgenium	
																		Cn 112 277 - - - Copernicium	
																		Uut 113 284 - - - Ununtrium	
																		Fl 114 289 - - - Flerovium	
																		Lv 116 282 - - - Livermorium	
																		Uus 117 293 - - - Ununseptium	
																		Uuo 118 294 - - - Ununoctium	



		6										7																
		Lanthanoids										Actinoids																
Ce 58	Pr 59	Nd 60	Pm 61	Sm 62	Eu 63	Gd 64	Tb 65	Dy 66	Ho 67	Er 68	Tm 69	Yb 70	Lu 71	Th 90	Pa 91	U 92	Np 93	Pu 94	Am 95	Cm 96	Bk 97	Cf 98	Es 99	Fm 100	Md 101	No 102	Lr 103	
140.116	140.90765	144.242	144.9127	150.36	151.964	157.25	168.92535	162.500	164.93032	167.259	168.93421	173.054	174.9668	1.12 4,3	1.13 3	1.14 3	1.13 3	1.17 3	1.2 3,2	1.2 3	1.2 3	1.23 3	1.23 3	1.23 3	1.23 3	1.25 3	1.1 3	1.27 3
Cerium	Praseodymium	Neodymium	Promethium	Samarium	Europium	Gadolinium	Terbium	Dysprosium	Holmium	Erbium	Thulium	Ytterbium	Lutetium	Thorium	Protactinium	Uranium	Neptunium	Plutonium	Americium	Curium	Berkelium	Californium	Einsteinium	Fermium	Mendelevium	Nobelium	Lawrencium	
232.03806	231.03688	238.02891	237.0482	244.0642	243.0614	247	247.0703	251.0796	252.03	257.0951	258.01	259.1009	260.1053	1.3 4	1.5 5	1.38 6	1.36 5	1.28 4	1.3 3	1.3 3	1.3 3	1.3 3	1.3 3	1.3 3	1.3 3	1.3 3	1.3 3	1.3 3

Atomic masses are measured relative to the carbon isotope ¹²C (IUPAC-2010)