


1. (a)[2] Graph the pair of functions  $\sqrt{x}$  and  $\ln x$  in the same coordinate system and identify the one that approaches  $\infty$  faster as  $x$  approaches  $\infty$ .



$x$	$\sqrt{x}$	$\ln x$
100	10	4.6
1000	31.6	6.9
$10^6$	1000	13.8
$\vdots$	$\vdots$	$\vdots$

↑  
FASTER

(b)[3] Find  $\lim_{x \rightarrow \infty} (\ln(2x^6 + 1) - 3 \ln x^2) = \lim_{x \rightarrow \infty} \ln \frac{2x^6 + 1}{x^6}$

$= \underline{\underline{\ln 2}}$

since  $\lim_{x \rightarrow \infty} \frac{2x^6 + 1}{x^6} = \lim_{x \rightarrow \infty} \frac{2x^6}{x^6} = 2$

2. True/false questions. Decide whether the statements in questions (a) - (b) are true or false (circle your choice). You must correctly justify your answer to receive credit.


(a)[2] Every function defined on a closed interval  $[a, b]$  has an absolute maximum and an absolute minimum.

must be continuous!

TRUE

**FALSE**

or, give an example:



defined on  $[a, b]$   
 but no max, no min

(b)[2] If  $f'(c) = 0$ , then  $f(x)$  must have an extreme value (i.e., either minimum or maximum) at  $c$ .

$f(x) = x^3$  satisfies  $f'(0) = 0$   
 but does not have extreme value at 0


TRUE

**FALSE**

(c)[2] The function below has two critical numbers (critical points).


TRUE

**FALSE**



has three critical points

3. Consider the function  $f(x)$  given in the graph below.



(a)[2] What is the value of  $f'(1.8)$ ? Explain your answer.


slope of the line =  $\frac{-1-1}{2-1} = -2$

(b)[2] The function  $f(x)$  has a corner (cusp) at  $x = 2$ . This means that  $f'(2)$  does not exist. Explain why.

as  $x \rightarrow 2^-$  slopes (are constant, equal to  $-2$ ) approach  $-2$   
 as  $x \rightarrow 2^+$  slopes approach  $0$  so slope at 2 d.n.e.

(c)[3] Sketch the graph of  $f'(x)$ .

$f'(x) = 0$  on  $(-\infty, 1)$        $f'$  positive, decreasing on  $(3, \infty)$   
 $f'(x) = -2$  on  $(1, 2)$   
 $f'(x) = 0$  on  $(2, 3)$        $f'$  d.n.e. at  $1, 2, 3$



4. Consider the discrete-time dynamical system  $p_{t+1} = 1.4p_t(1-p_t)$ , where  $p_t$  is a population of mice in thousands.

(a)[2] Find all equilibrium points of the system.

$$p^* = 1.4p^*(1-p^*) \rightarrow p^*(1-1.4(1-p^*)) = 0$$

$$\text{so } \underline{\underline{p^* = 0}} \quad \text{or} \quad 1-1.4+1.4p^* = 0$$

$$1.4p^* = 0.4, \quad p^* = \frac{0.4}{1.4} = \underline{\underline{\frac{2}{7}}}$$

(b)[3] Determine whether each of the equilibrium points you found in (a) is stable or unstable.

$$f(x) = 1.4x(1-x) = 1.4x - 1.4x^2$$

$$f'(x) = 1.4 - 2.8x$$

$$f'(0) = 1.4 \dots \text{ since } |f'(0)| > 1, \quad 0 \text{ is } \underline{\underline{\text{unstable}}}$$

$$f'\left(\frac{2}{7}\right) = 1.4 - 2.8 \cdot \frac{2}{7} = 0.6 = \frac{3}{5}$$

$$|f'\left(\frac{2}{7}\right)| < 1, \quad \frac{2}{7} \text{ is } \underline{\underline{\text{stable}}}$$

(c)[2] Explain what your answer to (b) means for the population of mice.

populations near 0 (ie small population) will move away from it, ie will increase

populations near  $\frac{2}{7}$  thousands will move closer (ie increase or decrease) to  $\frac{2}{7}$  thousands

Continued on next page

5. (a)[1] What is the per capita production in the context of population models?

number of offspring produced by  
a single individual

(b)[1] Identify per capita production in the population discrete system  $p_{t+1} = \frac{2p_t}{1 + 0.005p_t}$ .

$$\frac{2}{1 + 0.005p_t}$$

(c)[3] Find all equilibrium points of the system in (b).

$$p^* = \frac{2p^*}{1 + 0.005p^*} \rightarrow p^* \left( 1 - \frac{2}{1 + 0.005p^*} \right) = 0$$

So  $p^* = 0$  or  $1 + 0.005p^* = 2 \rightarrow p^* = \frac{1}{0.005} = \underline{\underline{200}}$

(d)[2] Consider the largest equilibrium point that you found in (c). Is it stable or not?

$$f(x) = \frac{2x}{1 + 0.005x}$$

$$f'(x) = \frac{2(1 + 0.005x) - 2x(0.005)}{(1 + 0.005x)^2} = \frac{2}{(1 + 0.005x)^2}$$

$$f'(200) = \frac{2}{(1 + (0.005)(200))^2} = \frac{1}{2}$$

↑  
Stable

Since  $|f'(200)| < 1$

Continued on next page

6. (a) [2] Give the statement of the Extreme Value Theorem.

If  $f$  is continuous on  $[a, b]$ ,  $[a, b]$  closed

Then  $f$  has an abs. max. and an abs. min. in  $[a, b]$

(b) [3] Find the absolute maximum and the absolute minimum of the function  $f(x) = \frac{\ln x}{x^2}$  on the interval  $[1, 4]$ .

$$f'(x) = \frac{\frac{1}{x} \cdot x^2 - (\ln x)(2x)}{(x^2)^2} = \frac{x(1 - 2 \ln x)}{x^4}$$

$$= \frac{1 - 2 \ln x}{x^3}$$

$$f'(x) = 0 \dots 1 - 2 \ln x = 0, \ln x = \frac{1}{2}, x = e^{1/2} = \sqrt{e} \approx 1.65$$

$f'(x)$  dne... no such points in  $[1, 4]$

$x$	$f(x) = \frac{\ln x}{x^2}$
$\sqrt{e}$	$\frac{\ln \sqrt{e}}{(\sqrt{e})^2} = \frac{1/2}{e} = \frac{1}{2e} \approx 0.18$
1	0
4	$\frac{\ln 4}{16} \approx 0.09$

abs min = 0, at  $x = 1$   
 abs max =  $\frac{1}{2e}$  at  $x = \sqrt{e}$   
 $\uparrow$   $\uparrow$   
 0.18 1.65

Continued on next page

7. [3] Find all critical numbers (critical points) of the function  $f(x) = (x - 1)^{2/3} + 1$ .

$$f'(x) = \frac{2}{3} (x-1)^{-\frac{1}{3}} = \frac{2}{\sqrt[3]{x-1}}$$

$$f'(x) = 0 \dots \text{no } x$$

$$f'(x) \text{ dne when } x = 1$$

Since  $x = 1$  is in domain of  $f(x)$ , it is a c.p.

so  $x = 1$  is (the only) c.p.