


1. (a)[2] Find all solutions of the equation  $\cos x = -1/2$ .




Know:  $\cos \frac{\pi}{3} = \frac{1}{2}$

$$x_1 = \pi - \frac{\pi}{3} = \frac{2\pi}{3} + 2\pi k$$

$$x_2 = \pi + \frac{\pi}{3} = \frac{4\pi}{3} + 2\pi k$$

- (b)[3] A population  $P(t)$  of brown wolves in a small region of southern Nunavut has been known to fluctuate between the low of 120 wolves and high of 400 wolves, with a period of 14 years (i.e., increases from 120 to 400 and decreases back to 120 in 14 years). Use a trigonometric function to find a formula for  $P(t)$  as function of time  $t$ , measured in years.



Use cosine

period  $\frac{2\pi}{a} = 14 \rightarrow a = \frac{\pi}{7}$

$\cos\left(\frac{\pi}{7}x\right) \rightarrow$  adjust for min/max

$$-140 \cos\left(\frac{\pi}{7}x\right) + 260$$

$$\text{so } P(t) = 260 - 140 \cos\left(\frac{\pi}{7}t\right)$$


2. (a)[2] Is the formula  $\arcsin(\sin x) = x$  true for all real numbers  $x$ ? Justify your answer.

no, only true for  $x$  in  $[-\frac{\pi}{2}, \frac{\pi}{2}]$

for example,  $\arcsin(\sin \pi) = \arcsin(0) = 0 \neq \pi$

(b)[2] Find the domain of the function  $f(x) = \arcsin(1/x)$ . Explain your answer.

GRAPH



$\frac{1}{x} \dots x \neq 0, \arcsin \dots -1 \leq \frac{1}{x} \leq 1$


OR ALGEBRA

$\frac{1}{x} \leq 1 \rightarrow x \geq 1 \dots [1, \infty)$  and

$\frac{1}{x} \geq -1 \dots x \leq -1 \dots (-\infty, -1]$

$(-\infty, -1]$  and  $[1, \infty)$

(c)[2] Find the range of the function  $y = \arcsin(x^2)$ . Explain your answer.




$x^2 \geq 0$   
 so  $\arcsin(x^2) \geq 0$

range =  $[0, \frac{\pi}{2}]$

3. Find the following limits

$$\begin{aligned}
 \text{(a)[3]} \quad \lim_{x \rightarrow 4} \frac{\frac{1}{\sqrt{x}} - \frac{1}{2}}{x-4} &= \frac{0}{0} \\
 &= \lim_{x \rightarrow 4} \frac{\frac{2-\sqrt{x}}{2\sqrt{x}}}{x-4} = \lim_{x \rightarrow 4} \frac{2-\sqrt{x}}{2\sqrt{x}} \cdot \frac{1}{x-4} \\
 &= \lim_{x \rightarrow 4} \frac{2-\sqrt{x}}{2\sqrt{x}(\sqrt{x}-2)(\sqrt{x}+2)} = \\
 &= \lim_{x \rightarrow 4} \frac{-1}{2\sqrt{x}(\sqrt{x}+2)} = -\frac{1}{16}
 \end{aligned}$$

$$\text{(b)[2]} \quad \text{Find } \lim_{x \rightarrow (\pi/2)^+} x^2 \sec x = x^2 \cdot \frac{1}{\cos x} = \left(\frac{\pi^2}{2}\right) \cdot \frac{1}{0^-} = -\infty$$



Continued on next page

4. Consider the function

$$f(x) = \begin{cases} \frac{x-1}{x^3-x} & \text{if } x \neq 1 \\ 1/2 & \text{if } x = 1 \end{cases}$$

(a)[2] Is  $f(x)$  continuous at  $x = -2$ ? Explain why or why not.

near  $x = -2 \dots f(x) = \frac{x-1}{x^3-x}$  is continuous  
 as quotient of continuous functions with  
 denominator  $(-2)^3 - (-2) = -6 \neq 0$

(b)[3] Is  $f(x)$  continuous at  $x = 1$ ? Explain why or why not.

$$\lim_{x \rightarrow 1} f(x) = \lim_{x \rightarrow 1} \frac{\cancel{x-1}}{x(\cancel{x-1})(x+1)} = \frac{1}{2} = f(1)$$


YES

(c)[2] Find all real numbers  $x$  where the function  $f(x) = \sqrt{\tan x}$  is continuous.

$$\Rightarrow \tan x \geq 0$$

$$x \text{ is in } [0, \frac{\pi}{2})$$

$$\text{so } [0 + \pi k, \frac{\pi}{2} + \pi k)$$



Continued on next page

5. Let  $f(x) = \sqrt{x}$ .

(a)[1] Find the average rate of change of  $f(x)$  on  $[4, 4.2]$ .

$$\frac{f(4.2) - f(4)}{0.2} = 0.2469$$

(b)[1] Find the average rate of change of  $f(x)$  on  $[4, 4.1]$ .

$$\frac{f(4.1) - f(4)}{0.1} = 0.2485$$

(c)[3] What number are the values in (a) and (b) supposed to approach? Explain why they approach that number and not some other number.

$$f'(x) = \frac{1}{2\sqrt{x}} \quad f'(4) = \frac{1}{2\sqrt{4}} = \frac{1}{4}$$

they are supposed to approach  $\frac{1}{4} = 0.25$

since average rate of change approaches  
instantaneous rate of change as  
interval shrinks to a point

6. (a)[2] Find  $f'(x)$ , if  $f(x) = 3^{\ln x} + (\ln x)^3 + (\ln 3)^3$ .

$$f'(x) = 3^{\ln x} \ln 3 \cdot \frac{1}{x} + 3(\ln x)^2 \cdot \frac{1}{x} + 0$$

(b) [2] Find  $f'(0)$ , if  $f(x) = \arcsin x + (\arcsin x)^2$ .

$$f'(x) = \frac{1}{\sqrt{1-x^2}} + 2\arcsin x \cdot \frac{1}{\sqrt{1-x^2}}$$

$$f'(0) = \frac{1}{\sqrt{1}} + \underbrace{2\arcsin 0}_0 \cdot \frac{1}{\sqrt{1}} = \underline{\underline{1}}$$

(c)[3] Find the equation of the line tangent to the graph of  $y = \frac{e^x + 1}{x}$  at the point where  $x = 1$ .

$$x = 1 \rightarrow y = e + 1$$

$$y' = \frac{e^x \cdot x - (e^x + 1) \cdot 1}{x^2} \quad \therefore y'(1) = \frac{e - (e + 1)}{1} = -1$$

$$\text{so } y - (e + 1) = -1(x - 1)$$


$$\underline{\underline{y = -x + e + 2}}$$

7. (a)[2] Let  $g(x) = x^2\sqrt{f(x)}$ , where  $f$  is a differentiable function such that  $f(1) = 4$  and  $f'(1) = 1$ . Find  $g'(1)$ .

$$g'(x) = 2x\sqrt{f(x)} + x^2 \cdot \frac{1}{2} (f(x))^{-1/2} \cdot f'(x)$$

$$g'(1) = \underbrace{2 \cdot 1 \cdot \sqrt{4}}_4 + 1 \cdot \frac{1}{2} \cdot \frac{1}{\sqrt{4}} \cdot 1 = 4 + \frac{1}{4} = \frac{17}{4}$$

(b)[3] The graphs of the functions  $f(x)$  and  $g(x)$  are given below. Compute  $(fg)'(3)$ , i.e. compute the derivative of the product of  $f(x)$  and  $g(x)$  when  $x = 3$ .



$$(fg)'(3)$$

$$= f'(3) \cdot g(3) + f(3) \cdot g'(3)$$

$$= \frac{1}{2} \cdot 2 + 4 \cdot (-1)$$

$$= -3$$