

## MAT 1341A – Test 2 – DGD 1, 2015

19 October, 2015.

Instructor – Barry Jessup.

Family Name: \_\_\_\_\_

First Name: \_\_\_\_\_

Student number: \_\_\_\_\_

Your multiple choice answers →	{	1	
		2	
		3	
For the marker's use only →	{	4	
		5	
		6	
		[Bonus] 7	
		Total	

**PLEASE READ THESE INSTRUCTIONS CAREFULLY.**

1. Read each question carefully, and **answer all questions in the space provided after each question.** For questions 4 to 7, you may use the backs of pages if necessary, but be sure to indicate to the marker that you have done this.
2. Questions 1 to 3 are multiple choice. These questions are worth 1 point each and no part marks will be given. Please record your answers in the space provided above.
3. Questions 4 – 6 and are worth 6 points each, and part marks can be earned. **The correct answers here require justification written legibly and logically: you must convince the marker that you know why your solution is correct.**
4. Question 7 is a challenging bonus question and is worth 3 points. It is *much* more difficult to obtain marks in the bonus question, so spend your time accordingly. You can earn 100% without attempting Q.7.
5. Where it is possible to check your work, do so.
6. Good luck! Bonne chance!

1. Which of the following are subspaces of  $\mathbf{R}^3$ ?

$$U = \{(x, y, z) \in \mathbf{R}^3 \mid x - 2y + z = 0\}$$

$$V = \{(x, xy, y) \in \mathbf{R}^3 \mid x, y \in \mathbf{R}\}$$

$$W = \{(x, y, z) \in \mathbf{R}^3 \mid 2x - 5z = 0\}$$

$$X = \{(x + y, y, x - 2y) \mid x, y \in \mathbf{R}\}$$

- A. Only  $U$  and  $V$
- B. Only  $U$  and  $W$
- C. Only  $W$  and  $X$
- D. Only  $U$ ,  $V$  and  $W$
- E. Only  $U$ ,  $V$  and  $X$
- F. Only  $U$ ,  $W$  and  $X$

2. It is known that a subspace  $Y$  of  $\mathbf{R}^{110}$  can be spanned by 96 vectors, and that  $Y$  has a linearly independent set with 71 vectors. Then it is always true that:

- A.  $\dim Y < 71$
- B.  $\dim Y > 71$
- C.  $71 < \dim Y \leq 96$
- D.  $71 \leq \dim Y < 96$
- E.  $71 \leq \dim Y \leq 96$
- F. None of the above is true.

3. Suppose  $\{u, v\}$  is a linearly **independent** set in vector space  $V$ , and that  $w \in V$  is chosen so that  $\{u, v, w\}$  is linearly **independent**. Which of the following statements is **ALWAYS** true?

- A.  $\{u, w\}$  is linearly dependent.
- B.  $\{v, w\}$  is linearly dependent.
- C.  $\{u, v\}$  is linearly dependent.
- D.  $u \in \text{span}\{v, w\}$ .
- E.  $v \in \text{span}\{u, w\}$ .
- F.  $w \notin \text{span}\{u, v\}$ .

4. Recall the vector space  $\mathcal{P}_2 = \{a + bx + cx^2 \mid a, b, c \in \mathbf{R}\}$  of polynomial functions of degree at most 2, and define

$$W = \{p \in \mathcal{P}_2 \mid p(3) = 0\}.$$

- a) Show that  $W = \text{span}\{x - 3, x^2 - 3x\}$ . (*Hint: recall the Factor Theorem: if  $p$  is any polynomial and  $p(a) = 0$  for some  $a \in \mathbf{R}$ , then  $p(x) = (x - a)q(x)$  for some polynomial  $q$  of degree one less than that of  $p$ .*)
- b) Explain why  $W$  is a subspace of  $\mathcal{P}_2$  *without using the subspace test*.
- c) Find a basis for  $W$ . (You may use without proof the fact proved in class that  $\{1, x, x^2\}$  is linearly independent.)
- d) Find  $\dim W$ .

*(Remember that you must justify your answers.)*



5. Let  $\mathbf{M}_{22}$  denote the vector space of 2 by 2 matrices with real entries, and define

$$U = \left\{ \begin{bmatrix} a & b \\ a+b & c \end{bmatrix} \in \mathbf{M}_{22} \mid a, b, c \in \mathbf{R} \right\}.$$

a) Either check that  $U$  is closed under addition, or express  $U$  in another form so you can simply state a theorem that guarantees that  $U$  is a subspace.

(For parts (b) and (c) you may assume that  $U$  is a subspace of  $\mathbf{M}_{22}$ .)

b) Find a basis for  $U$ , and hence find  $\dim U$ .

c) Give a basis for  $U$  different from the one you gave in (b).

*(Remember that you must justify your answers.)*



6. State whether each of the following statements is (always) true, or is (possibly) false, in the box after the statement.

- If you say the statement may be false, you must give an explicit example - with numbers, matrices, or functions, as is appropriate!
- If you say the statement is always true, you must give a clear explanation.

a)  $X = \{f \in \mathbf{F}(\mathbf{R}) \mid f(x) \geq -1 \text{ for all } x \in \mathbf{R}\}$  is a subspace of  $\mathbf{F}(\mathbf{R}) = \{f \mid f : \mathbf{R} \rightarrow \mathbf{R}\}$ .

ANSWER

b) If  $V$  is a vector space and  $\{v_1, v_2, v_3\} \subset V$  is linearly independent, then  $\{v_1, v_2\}$  is also linearly independent.

ANSWER

**6 (cont.).**

c)  $\left\{ \begin{bmatrix} a & a \\ b & c \end{bmatrix} \in \mathbf{M}_{2,2} \mid a, b, c \in \mathbf{R} \right\}$  is a subspace of  $\mathbf{M}_{2,2}$  of dimension 3.

ANSWER

d) If  $v_1, v_2, v_3$  and  $v_4$  are non-zero vectors in a vector space  $V$ , and  $U = \text{span}\{v_1, v_2, v_3, v_4\}$  then  $\dim U = 4$ .

ANSWER

**7.** [Challenge/Bonus]

Suppose  $U$  and  $W$  are two 3-dimensional subspaces of  $\mathbf{R}^5$ .

Explain carefully why there is a non-zero vector in  $U \cap W = \{v \in \mathbf{R}^5 \mid v \in U \text{ and } v \in W\}$ .

*Hint: Assume that  $U \cap W = \{0\}$  and find a contradiction.*

**Note: You cannot choose  $U$  or  $W$ . Your explanation must work for all 3-dimensional subspaces  $U$  and  $W$  of  $\mathbf{R}^5$ .**

