

School of Mathematics and Statistics
Carleton University
Math. 1005A, Winter 2011
Mock TEST 6

Any non-programmable calculator permitted, 1 blank sheet permitted for roughs

Print Name :

Student Number:

Tutorial Section (A1, A4, ...):

PART I: Multiple Choice Questions

(Choose and CIRCLE only ONE answer - No part marks here.)

1. [2 marks] Which of the following statements is true about the Fourier series of the function f defined on $(-\pi, \pi)$ by $f(x) = -1, -\pi < x < 0$ and $f(x) = 1, 0 < x < \pi$?
 (a) $a_n = 0$ for all $n \geq 1$, (b) $b_2 = 0$, (c) $a_0 \neq 0$, (d) $b_n = 0$ for all $n \geq 1$.
2. [2 marks] Find the Fourier cosine series of the function f defined on $(0, \pi)$ by $f(x) = 1, 0 < x < 1$ and $f(x) = 0, 1 < x < \pi$.
 (a) $\frac{2}{\pi} \sum_{n=1}^{\infty} \frac{\sin n}{n} \cos nx$, (b) $\frac{1}{\pi} + \sum_{n=1}^{\infty} \frac{\sin n}{n} \cos nx$ (c) $\frac{1}{\pi} + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{\sin n}{n^2} \cos nx$ (d) $\frac{1}{\pi} + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{\sin n}{n} \cos nx$.
3. [2 marks] The first term term, b_1 , of the Fourier sine series of the function $f(x) = x^2$ for $0 < x < \pi$ is given by:
 (a) 0, (b) $\frac{8}{\pi} \sin x$, (c) $2\pi - \frac{8}{\pi} \sin x$, (d) $2\pi + \frac{8}{\pi} \sin x$.
4. [2 marks] Find the first term, a_0 , of the Fourier cosine series of the function $f(x) = \frac{\pi}{4} - \frac{x}{2}$ where $0 < x < \pi$.
 (a) $a_0 = \frac{2}{\pi}$, (b) $a_0 = 0$, (c) $a_0 = 1$, (d) $a_0 = \frac{\pi}{2}$.
5. [2 marks] Among the following pairs of functions (f, g) , (g, h) and (f, h) where

$$f(x) = \cos x, \quad g(x) = \sin^2 x, \quad h(x) = x^2,$$

which pairs represent functions that are orthogonal to each other on the interval $(0, \pi)$?

- (a) (f, h) but not (f, g) , (b) (f, g) but not (f, h) or (g, h) , (c) (g, h) but not (f, g) or (f, h) , (d) (f, h) but not (g, h) .

PART II: Show all work here and give details.

No additional pages will be accepted

6. [10 marks] Let $f(x) = \begin{cases} 1, & -1 \leq x \leq 0 \\ x+1, & 0 < x \leq 1 \end{cases}$. Find the Fourier series of f in the form

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} [a_n \cos(n\pi x) + b_n \sin(n\pi x)]$$

where $n \geq 1$.

7. [10 marks] Let $f(x) = x^2$ for $0 \leq x \leq 1$. Find the a_n in the half-range cosine series of f (which is of the form $\sum_{n=1}^{\infty} a_n \cos(n\pi x)$)