

CONCORDIA UNIVERSITY
Department of Mathematics & Statistics

Course	Number	Sections	
Mathematics	205	All	
Examination	Date	Pages	
Final	December 2013	2	
Instructors:	G. Bobos, J. Brody, A Iovita, H. Proppe	Course Examiners	A. Atoyan & H. Proppe
Special Instructions:	Only calculators approved by the Department are allowed		

MARKS

[10] 1. (a) Sketch the graph of the function

$$f(x) = \begin{cases} -\sqrt{9-x^2} & \text{if } -3 \leq x < 0 \\ 1 - |x-4| & \text{if } 0 \leq x \leq 5 \end{cases}$$

on the interval $-3 \leq x \leq 5$ and calculate the definite integral $\int_{-3}^5 f(x) dx$ as the signed area between the graph of f and the x -axis (do not antidifferentiate).

(b) Use the Fundamental Theorem of Calculus to find a function $f(x)$ and

$$\text{a number } a \text{ so that } a + \int_4^x \frac{f(t)}{t^2} dt = 2\sqrt{x} \text{ for all } x > 0.$$

[11] 2. Calculate the following indefinite integrals:

$$\text{(a)} \quad \int \frac{(x + x^{3/2})^2}{\sqrt{x}} dx \qquad \text{(b)} \quad \int \frac{5x - 3}{x^2 - 2x - 3} dx$$

[10] 3. Find the antiderivative $F(t)$ of the function $f(t)$ that satisfies the given condition:

$$\text{(a)} \quad f(t) = (t-1)(t^2-2t)^{10}, \quad F(1) = 0. \quad \text{(b)} \quad f(t) = \sin^3 t \cos^2 t, \quad F\left(\frac{\pi}{2}\right) = 10.$$

[12] 4. Evaluate the following definite integrals (give the exact answers):

$$\text{(a)} \quad \int_1^e x \ln^2 x dx \qquad \text{(b)} \quad \int_{-\pi/4}^{\pi/4} \tan^2 x dx$$

[8] 5. Evaluate the given improper integral or show that it diverges:

$$(a) \int_1^{\infty} \frac{1}{x(\ln x)^2} dx \quad (b) \int_{-1}^0 \frac{2}{x^2 - 1} dx$$

[17] 6. (a) Sketch the curves $y = 2 - x^2$ and $y = -x$ and find the area enclosed by these curves.

(b) Sketch the region enclosed by $f(x) = x^2$, the x -axis, the y -axis, and the line $x = 1$. Find the volume of the solid generated by revolving this region about the y -axis.

(c) Find the average value of the function $f(x) = \sqrt{25 - x^2}$ on the interval $[-5, 5]$.

[9] 7. Find the limit of the sequence $\{a_n\}$ or prove that the limit does not exist:

$$(a) a_n = \frac{\ln(n)}{\ln(2n)} \quad (b) a_n = \frac{n}{n+1} \cos(n\pi) \quad (c) a_n = n - \sqrt{n^2 - n}$$

[11] 8. Determine whether the series is divergent or convergent, and if convergent, then absolutely or conditionally :

$$(a) \sum_{n=1}^{\infty} \frac{\arctan n}{n^{1.2}} \quad (b) \sum_{n=1}^{\infty} (-1)^{n-1} e^{2/n} \quad (c) \sum_{n=1}^{\infty} \frac{n^5}{5^n}$$

[6] 9. Find the radius and interval of convergence of the power series $\sum_{n=1}^{\infty} \frac{(2x-1)^n}{n4^n}$

[6] 10. (a) Use the MacLaurin series for e^{-x} to find the MacLaurin series for $f(x) = e^{-x^2}$.

(b) Find an antiderivative $F(x)$ for $f(x) = e^{-x^2}$ expressed as a power series.

[5] **Bonus Question.** Prove that if $a_n > 0$ and $\sum_{n=1}^{\infty} a_n$ converges, then $\sum_{n=1}^{\infty} a_n^2$ also converges. However, if the condition $a_n > 0$ is removed, show that this is not necessarily true, i.e. $\sum_{n=1}^{\infty} a_n^2$ may diverge.

The present document and the contents thereof are the property and copyright of the professor(s) who prepared this exam at Concordia University. No part of the present document may be used for any purpose other than research or teaching purposes at Concordia University. Furthermore, no part of the present document may be sold, reproduced, republished or re-disseminated in any manner or form without the prior written permission of its owner and copyright holder.