

BIO 2135 - Animal Form and Function
Midterm examination
Worth either 10% or 15% of your final grade

Tuesday February 8, 2011

- a) Place your name and student number in the space provided below. Be sure that your name is on the top of each page because the exam will be separated to facilitate marking
- b) Circle the lab section for your lab.
- c) Check to be sure that your exam is complete with a total of 12 pages including this one
- d) Answer all questions in the space provided on the exam. Do not transfer answers to the back of the page
- e) The exam is out of 95 pts.

Name:

Student No:

Exams will be returned on the lab so please circle your lab section:

Thursday: A1-BSC312, A3-BSC330, A5-BSC335 A7-BSC310

Friday: A2-BSC312, A4-BSC330 A6-BSC335

•

Name: Student Number:

15 pts Part 1. Briefly explain what each of the following biological terms means. Where possible include an example in your definition from a group or an organism to which the term applies.

Statolith

Cnidocil

Spiral cleavage

Pellicle

Porocyte

Name: Student Number:

20 pts Part 2 Answer each of the following multiple choice questions by placing an X in the space to the left of the correct choice. There is only one correct answer for each question.

2.1 Where are the choanocytes in a syconoid sponge

- _____ A: Spongocoel
- _____ B: Incurrent canal
- _____ C: Radial canal
- _____ D: Choanocyte chambers
- _____ E: None of the above

2.2 A cnidarian skeleton is

- _____ A. Bony
- _____ B. Cartilagenous
- _____ C. Made of spicules
- _____ D. made of ossicles
- _____ E. hydrostatic

2.3 Name given to the stage in *Paramecium* conjugation when the two haploid micronuclei have fused to return to the diploid state.

- _____ A: Synkaryon
- _____ B: Prokaryon
- _____ C: Mesokaryon
- _____ D: Metakryon

2.4 This is the first type of cell that is attacked by the malaria parasite as it enters its human host.

- _____ A: Red Blood Cell
- _____ B: Liver Cell
- _____ C: Stomach cells
- _____ D: Salivary gland cell

2.5 The brain of a Cnidarian is best described as

- _____ A: nonexistent
- _____ B: enclosed in a protective cranium
- _____ C: as only a dense mass of nervous tissue surrounding the mouth
- _____ D: consisting of paired ganglia and paired nerve cord
- _____ E: Both b and d are correct

Name: Student Number:

2.6 The thin flat cells covering the outer surface of sponge

- _____ A. Choanocytes
- _____ B. Pinacocytes
- _____ C. Amebocytes
- _____ D. Sclerocytes
- _____ E. Archeocytes

2.7 The collar of a choanocyte

- _____ A. Regulates the diameter of the osculum
- _____ B. Filters food from the water
- _____ C. Regulates the diameter of the ostia
- _____ D. Secrete the digestive enzymes

2.8 A coelom is a body cavity lined with

- _____ A. Mesenteries
- _____ B. Epidermis
- _____ C. Mesodermal tissue
- _____ D. Ectodermal tissue
- _____ E. Endoderm

2.9 In the ciliates this structure serves as the genetic reserve of the cell.

- _____ A. Macronucleus
- _____ B. Polyploid nucleus
- _____ C. Triploid nucleus
- _____ D. Micronucleus

2.10 In the Anthozoa these ciliated tracts circulate water through the gastrovascular cavity to maintain the hydrostatic skeleton.

- _____ A. Siphonoglyphs
- _____ B. Mesenteries
- _____ C. Manubria
- _____ D. Rhopalia
- _____ E. Oral discs

2.11 The chills and fever of malaria are correlated with

- _____ A. cycles of shizogony in the liver cells
- _____ B. release of the sporozoites from the oocysts
- _____ C. escape of merozoites from the red blood cells
- _____ D. the formation of the gametocytes

Name: Student Number:

2.12 In protists, both gas exchange and excretion occur across the plasma membrane using

- _____ A. Endocytosis
- _____ B. Active transport
- _____ C. Simple diffusion
- _____ D. Exocytosis
- _____ E. Pinocytosis

2.13 In Cnidarians intracellular digestion occurs

- _____ A. in the gastrovascular cavity
- _____ B. in the nutritive muscular cells of the gastrodermis
- _____ C. in the mesohyl
- _____ D. in the epithelio-muscular cells of the gastrodermis
- _____ E. in the stomach

2.14 Correct sequence in a schyphozoan life cycle.

- _____ A: Strobila, Ephyra, Schyphostome, Medusa, Planula.
- _____ B: Strobila, Planula, Schyphostome, Medusa, Ephyra.
- _____ C: Strobila, Planula, Schyphostome, Medusa, Ephyra.
- _____ D: Strobila, Ephyra, Medusa, Planula, Schyphostome.
- _____ E: None of the above.

2.15 Most ciliophoran (ciliate) protists

- _____ A. have flagella
- _____ B. have a cytostome
- _____ C. are marine
- _____ D. are mainly parasitic

2.16 The cavity of cnidarians that serves as a digestive chamber, and for exchange of gases and metabolic wastes.

- _____ A: Intestine
- _____ B: Spongocoel
- _____ C: Gastrocoel
- _____ D. Gastrovascular cavity
- _____ E. Archenteron

Name: Student Number:

2.17 Eggs in sponges are commonly formed from

- _____ A. Pinacocytes
- _____ B. Porocytes
- _____ C. Choanocytes
- _____ D. Sclerocytes
- _____ E. None of the above

2.18 The nematocyst is an organelle found in this type of Cnidarian cell.

- _____ A. Cnidocyte
- _____ B. Choanocyte
- _____ C. Epitheliomuscular cell
- _____ D. Nutritive muscular cell
- _____ E. None of the above

2.19 This membranous shelf inside the bell margin of hydrozoans is important for locomotion and unique to the class.

- _____ A. Statoblast
- _____ B. Hydranth
- _____ C. Coenosarc
- _____ D. Velum
- _____ E. Manubrium

2.20 In protists this form of reproduction involves the production of multiple nuclei by mitosis and after all the nuclei are produced cytoplasmic cell division separates each nucleus into a new cell.

- _____ A. Binary fission
- _____ B. Fragmentation
- _____ C. Budding
- _____ D. Conjugation
- _____ E. Multiple fission

Name: Student Number:

20 pts Part 3: Complete the following sentences using the appropriate terms. Place the term in the space in the sentence or at the end of the sentence.

- 3.1 Euglenozoans use this/these for locomotion. _____
- 3.2 Embedded in the epidermis, these protect a free-living flatworm from predation. _____
- 3.3 Anthozoans are missing this stage of the typical cnidarian life cycle. _____
- 3.4 This is the structure that keeps a tapeworm anchored in the intestine. _____
- 3.5 The dual roles of cells lining the cnidarian gastrovascular is reflected in their name. _____
- 3.6 Instead of dorsal and ventral sides, cnidarians have oral and this surface. _____
- 3.7 The junction of the oviduct, the opening of the seminal receptacle, and where the egg is usually fertilized in a parasitic flatworm. _____
- 3.8 During the evolution of the different animal phyla, this tissue is found for the first time in all the Bilateria. _____
- 3.9 Flatworms are hermaphrodites; this is the more scientific term or synonym used to indicate this. _____
- 3.10 This "cap" is found at the advancing edge of a pseudopod. _____
- 3.11 These structures found on the sides of a free-living flatworm's head are chemosensory. _____
- 3.12 The nematocyst is this type of subcellular structure. _____
- 3.13 Sporogony increases the number of malarial parasites in this host. _____
- 3.14 A flatworm's incomplete gut is missing this structure. _____
- 3.15 Initially, the pH of the digestive fluid in the food vacuole is this. _____
- 3.16 These polyps in a hydrozoan colony are responsible for reproduction. _____
- 3.17 When comparing the length of undulipods flagella are _____.
- 3.18 These delicate, minute anthozoans create a unique and highly productive marine environment. _____
- 3.19 The common name for the phylum Platyhelminthes. _____
- 3.20 In the deepest ocean depths, the shells of protists can only be made of this mineral. _____

Name: Student Number:

10 pts Part 4: Indicate whether the following statements are True or False by placing a T for true and an F for False in the space next to the question.

- _____ 4.1 Dyenin arms connect the out microtubular doublets to the central core of an undulipod.
- _____ 4.2 The egg forms from the microgametocyte while in the blood of the infected human host
- _____ 4.3 The outer layer of the tapeworm tegument is formed from syncytial epidermal cells
- _____ 4.4 Nutrients for the eggs in Platyhelminthes are produced in the vitelline glands.
- _____ 4.5 Each proglottid in a tapeworm can be considered an individual and the whole tapeworm a type of colony.
- _____ 4.6 Freshwater sponges remove excess water from their cells using water expulsion vesicles.
- _____ 4.7 The trochophore larva is an autapomorphy for the Platyzoa.
- _____ 4.8 The syconoid architecture is the most common sponge body plan.
- _____ 4.9 Spongin is made of protein that resembles collagen.
- _____ 4.10 Choanocytes are located in the choanocyte chambers of the leuconoid type of sponge.

Part 5 of the exam starts on the next page

Name: Student Number:

30 pts Part 5: Answer 5 of the following 8 questions in the space provided each answer is worth 6 points. If you answer more than five only the first five will be counted.

5.1 Briefly describe what a schizocoel is and how it is formed.

5.2 What are the different stages and their sequence in the fluke life cycle? Give the functional importance of three of them.

Name: Student Number:

5.3 Anthozoans such as *Metridium* (the sea anemones) have complete and incomplete septa; what are they and how do they differ from each other?

5.4 What is a rhopalium, what does it do and in what animals would you find it?

Name: Student Number:

5.5 Who are the Protostomia and what are the autapomorphies that define the group, explain two of them.

5.6 How is the nervous and sensory system organized in a free living flatworm like *Dugesia* (Planaria)

Name: Student Number:

5.7 What is the differences and similarities between mesohyl and mesoglea, give an example of an animal where you would find each?

5.8 Describe briefly how protozoans ingest their food.