



Examination Cover Sheet

COURSE: Chem	NUMBER: 205	SECTION(S): all	
EXAMINATION: <input checked="" type="checkbox"/> FINAL <input type="checkbox"/> ALTERNATE <input type="checkbox"/> DEFERRED VERSION: _____	DATE: Dec. 14th, 2015	TIME: 7-10 pm	PAGES: 7 Including cover DOUBLE-SIDED = 14 pages
INSTRUCTOR(S): C. Rogers, G. Dénès		DIVISION:	
MATERIALS ALLOWED: <input type="checkbox"/> No! Booklets <input checked="" type="checkbox"/> IBM (Scantron) <input checked="" type="checkbox"/> Blue or <input type="checkbox"/> Green <input checked="" type="checkbox"/> Printed Translation Dictionary Other NO _____ <input checked="" type="checkbox"/> Calculator <input type="checkbox"/> ENCS Approved <input type="checkbox"/> Other <input type="checkbox"/> No _____		INSTRUCTIONS: <input checked="" type="checkbox"/> Return all <input checked="" type="checkbox"/> Answer on Exam <input type="checkbox"/> Open book <input type="checkbox"/> Crib sheet Details _____	

Please print your name and I.D. number in the spaces below.

LAST NAME: _____

FIRST NAME: _____

STUDENT I.D. # _____

SPECIAL INSTRUCTIONS:


- Part I: Use the question pages for rough work; only the Scantron sheet will be graded.
- Parts II-III: Write your answers in the spaces provided (p.9-10). If you need extra space, use the blank pages (p.11-12). Please clearly indicate if those pages should be graded.
- Potentially useful information and a periodic table are on p.13-14.

Please leave this area blank:	Mark breakdown (professor use):
	Part I: / 76
	Part II: / 10
	Part III: / 14
	TOTAL: / 100

Part I – MCQs: The following 38 questions are multiple choice questions worth 2 marks each. There is ONE correct choice per question. You can do rough work on the exam paper, but it will not be marked. Enter your name, student ID number and MCQ answers on the Scantron sheet, in PENCIL. (Circles must be coloured in completely. Your birth date and other details are not needed.)

- All atoms of the same element have the same number of _____ in their nucleus. What is the missing word or words?
 - neutrons
 - electrons
 - protons
 - neutrons and protons
 - neutrons, protons, and electrons
- What method is used to determine the exact masses of isotopes and their relative abundances?
 - density measurements
 - filtration
 - electron microscopy
 - titration
 - mass spectrometry
- Which pair of atoms has the same number of neutrons?
 - $^{28}_{12}\text{Mg}$ and $^{28}_{11}\text{Na}$
 - $^{26}_{12}\text{Mg}$ and $^{28}_{12}\text{Mg}$
 - $^{28}_{14}\text{Si}$ and $^{28}_{12}\text{Mg}$
 - $^{28}_{14}\text{Si}$ and $^{28}_{11}\text{Na}$
 - $^{28}_{14}\text{Si}$ and $^{26}_{12}\text{Mg}$
- What are the values for x and y for $\text{Al}_x(\text{CO}_3)_y$?
 - x = 1 and y = 2
 - x = 2 and y = 3
 - x = 1 and y = 3
 - x = 3 and y = 1
 - x = 3 and y = 2
- The density of carbon tetrachloride is 1.59 g/mL. What is the mass of 50.00 mL of CCl_4 ?
 - 0.0318 g
 - 79.50 g
 - 79.5 g
 - 31.45 g
 - 31.4 g

6. Methane reacts with oxygen to form carbon dioxide and water.




Which one statement concerning this reaction is INCORRECT ?

- two moles of water are formed for one mole of methane consumed.
 - two molecules of oxygen are consumed per one molecule of methane consumed.
 - the combined mass of reactants consumed equals the mass of products formed.
 - two grams of oxygen are consumed for each gram of carbon dioxide formed.
 - three moles of gases are formed for each three moles of gases consumed.
7. What is the formula of chromium (III) oxide?
- Cr_2O_3
 - Cr_3O_2
 - CrO_3
 - Cr_3O
 - Cr_3O_3
8. What is the net ionic equation for the reaction of KOH and HCl in aqueous solution?
- $\text{H}^+(\text{aq}) + \text{KOH}(\text{aq}) \rightarrow \text{H}_2\text{O}(\ell) + \text{K}^+(\text{aq})$
 - $\text{K}^+(\text{aq}) + \text{Cl}^-(\text{aq}) \rightarrow \text{KCl}(\text{aq})$
 - $\text{HCl}(\text{aq}) + \text{KOH}(\text{aq}) \rightarrow \text{H}_2\text{O}(\ell)$
 - $\text{H}^+(\text{aq}) + \text{OH}^-(\text{aq}) \rightarrow \text{H}_2\text{O}(\ell)$
 - $\text{KOH}(\text{aq}) + \text{H}_2\text{O}(\ell) \rightarrow \text{H}^+(\text{aq}) + \text{K}(\text{OH})_2(\text{s})$
9. You measure water in two containers: a 10-mL graduated cylinder with marks at every mL, and a 1-mL pipet marked at every 0.1 mL. If you add the two measured volumes together in a calculation in your notebook, to what decimal place should you express the total?
- 0.001 mL
 - 0.01 mL
 - 0.1 mL
 - 1 mL
 - 10 mL
10. Which of the following is NOT a physical change?
- freezing a liquid to form a solid
 - dissolving a solid in water to form a solution
 - evaporating liquid water to form steam
 - burning a piece of paper to form gases and smoke
 - sublimating a piece of ice to form water vapour
11. What is the pH and description of a dilute HCl solution with a concentration of 0.030 M?
- pH 12.48, acidic solution
 - pH 12.48, basic solution
 - pH 7.00, neutral solution
 - pH 1.52, acidic solution
 - pH 1.52, basic solution


12. What is the energy of a single photon of γ -radiation with a wavelength of 1.33×10^{-12} m?
- 9.34×10^{-18} J
 - 4.41×10^{-15} J
 - 1.49×10^{-13} J
 - 8.34×10^{-12} J
 - 2.10×10^{-11} J
13. What type of orbital is designated by $n = 2, \ell = 0, m_\ell = 0$?
- $2s$
 - $2p$
 - $2d$
 - $2f$
 - None
14. Which of the following is associated with the value of the m_ℓ quantum number?
- orbital orientation
 - number of nodes
 - angular momentum quantum number
 - orbital size
 - de Broglie wavelength
15. How many electrons can be described by the quantum numbers $n = 4, \ell = 2, m_s = +1/2$?
- 1
 - 2
 - 3
 - 5
 - 10
16. What is the density of carbon dioxide gas at standard temperature and pressure (STP)?
- 0.89 g/L
 - 1.00 g/L
 - 1.32 g/L
 - 1.68 g/L
 - 1.96 g/L

17. The Boltzmann plot shown reveals the distribution of molecular speeds for 4 gases at the same temperature. Which choice is the most likely list of gases A, B, C, D to produce the curves shown on this graph?




- Xe, O₂, Ne, He
- Xe, He, Ne, O₂
- He, Ne, O₂, Xe
- He, O₂, Ne, Xe

18. Which type of elements have *least* affinity for electrons?
- transition metals
 - main group metals
 - noble gases
 - main group nonmetals
 - semiconductors
19. Which of the following elements is most likely to NOT obey the octet rule in a molecule?
- B
 - C
 - N
 - O
 - F
20. What is the correct Lewis structure for XeF_3^+ with the correctly placed formal charge?




21. A ball-and-stick structure of AsF_4^- is shown on the right. How is the *electron-pair* (basic) geometry described?
- pentagonal
 - seesaw
 - trigonal bipyramidal
 - tetrahedral
 - square planar




22. How is the *molecular* geometry of AsF_4^- (see figure in Q.21) described?
- pentagonal
 - see-saw
 - trigonal bipyramidal
 - tetrahedral
 - square planar
23. In AsF_4^- (see figure in Q.21), the lone pair is observed to be in an equatorial site, not axial. Why is this geometry favoured?
- Arsenic atoms are bigger than fluorine.
 - Arsenic is less electronegative than fluorine.
 - The bulky lone pair makes fewest 90° contacts with neighbouring As–F bond-pairs.
 - Lone pairs prefer to have as many neighbours as possible, no matter what the angle.
 - The lone pair prefers the warmer temperatures near the equator.

24. SF_4 is isoelectronic and isostructural with AsF_4^- . How is it polarized (see figures below)?




- a. It is polarized with the net dipole along the axial direction, positive at the top.
 b. It is polarized with the net dipole along the axial direction, positive at the bottom.
 c. It is polarized with the net dipole aligned with the S – lone-pair axis.
 d. It is polarized with the net dipole perpendicular to the S – lone-pair axis.
 e. It is not polarized.
25. Which of the images would best describe helium gas?
- a. A
 b. B
 c. C
 d. all
 e. none



26. According to VSEPR theory, what is the molecular geometry of IF_5 ?
- a. tetrahedral
 b. trigonal bipyramidal
 c. see-saw
 d. square pyramidal
 e. octahedral
27. What is the hybridization of the central nitrogen in IF_5 ?
- a. sp
 b. sp^2
 c. sp^3
 d. sp^3d
 e. sp^3d^2
28. What is the number of sigma (σ) bonds and pi (π) bonds in a molecule of carbon dioxide?
- a. 4 σ and 0 π
 b. 3 σ and 1 π
 c. 2 σ and 2 π
 d. 1 σ and 3 π
 e. 0 σ and 4 π
29. You are given temperature readings for two locations on Earth: 29°C and 256 K. Which choice correctly ranks these temperatures?
- a. 29°C is warmer than 256 K
 b. 29°C is the same as 256 K
 c. 29°C is colder than 256 K

30. Which element has the following electron configuration?



- P
- S
- O
- N
- F

31. Which of the following chemical equations refers to the second ionization of Ca?


- $\text{Ca(s)} + 2\text{e}^- \rightarrow \text{Ca}^{2-}(\text{s})$
- $\text{Ca(g)} \rightarrow \text{Ca}^{2+}(\text{g}) + 2\text{e}^-$
- $\text{Ca}^{2+}(\text{g}) \rightarrow \text{Ca}^+(\text{g}) + \text{e}^-$
- $\text{Ca(s)} \rightarrow \text{Ca}^+(\text{s}) + \text{e}^-$
- $\text{Ca}^+(\text{g}) \rightarrow \text{Ca}^{2+}(\text{g}) + \text{e}^-$

32. Which of the following statements about electrons and atoms is INCORRECT?

- Nearly all of the mass of an atom is localized in its central core or nucleus.
- Electrons are best described as standing waves oscillating in the space around the nucleus.
- Electrons circle the nucleus in elliptical orbits determined by the atom's nuclear charge.
- In addition to being farthest from the nucleus, the electrons in the valence shell are also responsible for an element's chemical properties.
- When an atom absorbs an ultraviolet photon, an electron moves to a higher energy level.

33. An aqueous solution containing cations (shown as smaller light spheres in left box) is mixed with a solution containing polyatomic anions (simplified as larger dark spheres in right box), in the quantities shown. The diagram on the right illustrates the result obtained.

(Note: Water molecules and spectator counter-ions not shown.)



Which combinations of cation and anion from the lists below are compatible with the observed results? **Possible cations:** K^+ , Ni^{2+} , NH_4^+ , Mg^{2+} & **Anions:** OH^- , NO_3^- , ClO_4^- , SO_4^{2-}

- KOH , $\text{Ni}(\text{NO}_3)_2$, NH_4ClO_4 , MgSO_4
- $\text{Ni}(\text{OH})_2$, NiSO_4 , $\text{Ba}(\text{OH})_2$, MgSO_4
- KOH , KNO_3 , NH_4NO_3 , NH_4ClO_4
- $(\text{NH}_4)_2\text{SO}_4$, K_2SO_4
- $\text{Ni}(\text{OH})_2$, $\text{Mg}(\text{OH})_2$

THE REMAINING MULTIPLE-CHOICE QUESTIONS DEAL WITH THIS SCENARIO:


Sulphuric acid is listed in a catalog as having a concentration of 95-98% H₂SO₄ by weight. The label on a bottle in the stockroom shows the density as 1.85 g/mL. In order to determine the actual concentration of acid in the bottle, a student dilutes 5.00 mL of the concentrated acid to 500. mL. She then takes four 10.00 mL samples and titrates each sample with standardized 0.1790 M sodium hydroxide solution. The results (in order) were:

Sample "run"	1 st	2 nd	3 rd	4 th
Volume NaOH (mL)	~20.70	20.30	20.40	20.35

34. Based on the advertised 95-98% by weight composition listed in the catalog and the density of 1.85 g/mL, what is the *expected* molarity range of the *concentrated* sulphuric acid solution?
- 36 – 37 mol/L
 - 18 – 19 mol/L
 - 17 – 20 mol/L
 - 3.6 – 3.7 mol/L
 - 1.7 – 1.9 mol/L
35. Would you consider the titration results to be relatively precise? Why or why not?
- Not precise – the average deviation of the volumes from runs 2-4 was > 5%.
 - Not precise – the relative error of the volumes from runs 2-4 was > 5%.
 - Precise – the average deviation of the volumes from runs 2-4 was < 5%.
 - Precise – the relative error of the volumes from runs 2-4 was < 5%.
 - More than one of these answers is correct.
36. What type of reaction is involved in this titration?
- precipitation
 - gas-forming
 - acid-base
 - redox
 - both b & c
37. How many moles of sulphuric acid were consumed on average in titration runs 2-4?
- 7.285×10^{-3} mol
 - 3.643×10^{-3} mol
 - 1.821×10^{-3} mol
 - 5.685×10^{-2} mol
 - 1.137×10^{-1} mol
38. What is the *measured* molarity of the *concentrated* sulphuric acid from the stock bottle?
- 36.4 mol/L
 - 18.2 mol/L
 - 3.64 mol/L
 - 0.364 mol/L
 - 0.182 mol/L

Part II: The following questions require short answers on this exam paper in the space provided. Be sure to give explanations where required.

39. (10 marks total) The two pictures below show the *gas-phase* reaction between reactants AH_3 and X_2 to form products A_2 and XH_2 , where A = dark coloured atoms and X = large, light coloured atoms. The small, light coloured atoms are hydrogen, H.



- (a) (2 marks) Which reactant is limiting? How can you tell?
- (b) (2 marks) Write a balanced equation for the reaction.
- (c) (3 marks) The elements A and X are actually real elements to be found between Li and Ne in the periodic table. Identify the reactants AH_3 and X_2 and products A_2 and XH_2 , and briefly explain your reasoning.
- Reactants: $AH_3 =$ _____ and $X_2 =$ _____
- Products: $A_2 =$ _____ and $XH_2 =$ _____
- Reasoning:
- (d) (3 marks) In a real reaction, with the same *proportions* of reactants as shown in the left box, the total pressure of the reactant mixture was 1.00 atm. If the temperature and container volume are not changed, what should the pressure be in the container after the reaction? Explain briefly.

Part III: Answer the following questions with complete written answers on this exam paper. Use the blank pages (p.11-12) if you need more space. Be sure to provide adequate explanations or details to justify your answers where it is appropriate. No explanation - half-marks max!

40. (14 marks total) The following reaction can be used to prepare iodine in the laboratory:



(a) (2 marks) What are the names of the following two compounds?

MnO_2 _____ Na_2SO_4 : _____

(b) (2 marks) What is the oxidizing agent, and what has been oxidized?

Oxidizing Agent: _____ Oxidized: _____

(c) (2 marks) What is the reducing agent, and what has been reduced?

Reducing agent: _____ Reduced: _____

(d) (6 marks) What mass of iodine can be obtained if 20.0 g of NaI is mixed with 10.0 g of MnO_2 and treated with an excess of sulphuric acid?

(e) (2 marks) After the reaction, if 14.3 g of solid I_2 were actually isolated, what was the reaction's percent yield?

BLANK PAGE FOR ROUGH WORK OR EXTRA SPACE

BLANK PAGE FOR ROUGH WORK OR EXTRA SPACE

POTENTIALLY USEFUL INFORMATION

Atomic mass unit	$1 \text{ amu} = 1.66054 \times 10^{-27} \text{ kg}$
Avogadro's number	$N = 6.022 \times 10^{23} \text{ mol}^{-1}$
Definition of <i>Joule</i>	$1 \text{ J} = 1 \text{ kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$
Definition of <i>Pascal</i>	$1 \text{ Pa} = 1 \text{ kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$
Gas constant	$R = 0.08206 \text{ L} \cdot \text{atm} \cdot \text{mol}^{-1} \text{K}^{-1} = 8.314 \text{ J} \cdot \text{mol}^{-1} \text{K}^{-1}$
Planck's constant	$h = 6.626 \times 10^{-34} \text{ J} \cdot \text{s}$
Pressure units	$760 \text{ mm Hg} = 760 \text{ torr} = 1 \text{ atm} = 101.325 \text{ kPa} = 1.01325 \text{ bar}$
Rydberg constant	$R = 1.0974 \times 10^7 \text{ m}^{-1}$
Speed of light	$c = 2.9979 \times 10^8 \text{ m} \cdot \text{s}^{-1}$

Periodic Table of the Elements

1 H 1.008																	2 He 4.00	
3 Li 6.941	4 Be 9.012															9 F 18.998	10 Ne 20.18	
11 Na 22.99	12 Mg 24.31															17 Cl 35.45	18 Ar 39.95	
19 K 39.10	20 Ca 40.08	3 Sc 44.96	4 Ti 47.87	5 V 50.94	6 Cr 52.00	7 Mn 54.94	8 Fe 55.85	9 Co 58.93	10 Ni 58.69	11 Cu 63.55	12 Zn 65.39	13 B 10.81	14 C 12.01	15 N 14.007	16 O 15.999	35 Br 79.90	36 Kr 83.80	
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc (97.91)	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.41	49 In 114.82	50 Sn 118.71	51 Sb 121.76	52 Te 127.60	53 I 126.90	54 Xe 131.29	
55 Cs 132.91	56 Ba 137.33	La-Lu	72 Hf 178.49	73 Ta 180.95	74 W 183.84	75 Re 186.21	76 Os 190.2	77 Ir 192.22	78 Pt 195.08	79 Au 196.97	80 Hg 200.59	81 Tl 204.38	82 Pb 207.2	83 Bi 208.98	84 Po 208.98	85 At 209.99	86 Rn 222.02	
87 Fr 223	88 Ra 226.03	Ac-Lr	104 Rf (261)	105 Db (262)	106 Sg (263)	107 Bh (262)	108 Hs (265)	109 Mt (266)										

57 La 138.91	58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm (145)	62 Sm 150.35	63 Eu 151.97	64 Gd 157.25	65 Tb 158.93	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.93	70 Yb 173.04	71 Lu 174.97
89 Ac 227.03	90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np (237)	94 Pu (245)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (260)