

Sol. Attached.


CONCORDIA UNIVERSITY
Department of Mathematics & Statistics

Course	Number	Sections
Mathematics	201	All
Examination	Date	Duration
Midterm	23 October, 2016	1 h 30 min
Special	Only approved calculators are allowed	
Instructions:	Show all your work for full marks	

- [12] (a) Find the midpoint M of the line segment joining $A(-3, 1)$ and $B(3, -3)$, and then find the distance between M and the point A .
(b) Find the equation of the line that passes through the point $(8, -7)$ and is perpendicular to the line $4x = 20 + 2y$.
(c) Find the coordinates of the center and the radius of the circle whose equation is $x(4 - x) = y(y + 3)$.
- [9] Consider the quadratic function $f(x) = 3x^2 - 9x + 12$.
(a) Express $f(x)$ in the vertex form.
(b) Find the coordinates of the vertex and indicate whether it corresponds to the maximum or minimum of $f(x)$.
(c) Find the x - and y -intercepts of the graph $y = f(x)$.
- [5] Consider the functions $f(x) = \sqrt{4x + 3}$ and $g(x) = x^2 - 1$. Find $(f \circ g)$ and $(g \circ f)$, and determine the domain of $(f \circ g)$ and the domain of $(g \circ f)$.
- [6] Given the rational function $f(x) = \frac{x^4 - 16}{3(x^2 + 1)(x^2 - x - 6)}$, find:
(a) the x - and y -intercepts,
(b) all vertical asymptotes, if any,
(c) all horizontal asymptotes, if any.
- [12] Find the solutions of the following equations:
(a) $9^x = 3^x + 12$
(b) $\log_2 x + \log_2(x - 14) = 5$
(c) $5^{\log_5(x^2)} - 4 \cdot 3^{\log_3(x+1)} = 8$
- [6] Consider the function $f(x) = \ln(x + 3) + 6$.
(a) Find the inverse function $f^{-1}(x)$.
(b) Find the domain and range of $f(x)$ and the domain and range of $f^{-1}(x)$.

Bonus. [3]: Let $f(x)$ be a function defined for all real x . (a) If we now that the range of f is $(-2, 1)$, what is the range of $F(x) = [f(x)]^2$? And (b) explain why this information is sufficient to claim that $F(x)$ cannot be invertible function even if $f(x)$ is invertible.


Math 201 Mid Term Oct 2016



(i) Mid pt AB $\left\{ \begin{array}{l} x = \frac{x_1+x_2}{2} = \frac{-3+3}{2} = \frac{0}{2} = 0 \\ y = \frac{y_1+y_2}{2} = \frac{1+(-3)}{2} = \frac{-2}{2} = -1 \end{array} \right\} \Rightarrow \text{Mid pt is } M(0, -1)$

(ii) dist. MA = $\sqrt{(x_2-x_1)^2 + (y_2-y_1)^2}$
 $= \sqrt{(-3-0)^2 + (1-[-1])^2}$
 $= \sqrt{9+4} = \sqrt{13} = 5$

④



Step 1 Get slope of Required line

(i) slope of given line $4x = 20 + 2y$
 $4x - 20 = 2y$
 $y = 2x - 10$
 $\Rightarrow \text{slope} = 2 \text{ or } \frac{2}{1}$

(ii) Slope of \perp line is $-\frac{1}{2}$

Step 2 let $P(x, y)$ be Any pt. on Required line

Slope AB = $\frac{y_2-y_1}{x_2-x_1}$

$-\frac{1}{2} = \frac{y-(-7)}{x-8}$

$2(y+7) = -1(x-8)$

$2y+14 = -x+8$

$2y = -x-6 \Rightarrow y = -\frac{1}{2}x-3$

④

c) Step 1 $x(4-x) = y(y+3) \Rightarrow 4x-x^2 = y^2+3y \Rightarrow -x^2-y^2+4x+3y=0$
 (Get into Standard form) $\Rightarrow x^2+y^2-4x+3y=0$

Step 2
 (Complete square)

$x^2-4x+4 + y^2+3y+\frac{9}{4} = 0+4+\frac{9}{4}$

$(x-2)(x-2) + (y+\frac{3}{2})(y+\frac{3}{2}) = 4+\frac{9}{4}$

$(x-2)^2 + (y+\frac{3}{2})^2 = \frac{25}{4}$

$\sqrt{(x-2)^2 + (y+\frac{3}{2})^2} = \sqrt{\frac{25}{4}} = \frac{5}{2}$

\Rightarrow Centre is $(2, -\frac{3}{2})$, Radius = $\frac{5}{2}$

④

2. a)

$$y = 3x^2 - 9x + 12$$

$$y = 3\left(x^2 - 3x + \frac{9}{4}\right) + 12 = 3\left(\frac{9}{4}\right)$$

(3)

$$y = 3\left(x - \frac{3}{2}\right)\left(x - \frac{3}{2}\right) + 12 = \frac{21}{4}$$

$$y = 3\left(x - \frac{3}{2}\right)^2 + \frac{21}{4} \Rightarrow$$

b)

Vertex is $\left(\frac{3}{2}, \frac{21}{4}\right)$

opens up (concave up)

Vertex is a Minimum.

c)

x int

y int

$$y = 3x^2 - 9x + 12$$

$$3x^2 - 9x + 12 = 0$$

$$x^2 - 3x + 4 = 0$$

$$x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(4)}}{2(1)}$$


$$x = \frac{3 \pm \sqrt{-7}}{2}$$

No x intercept (or see graph)

$$y = 3x^2 - 9x + 12$$

$$y = 3(0)^2 - 9(0) + 12$$

$$\Rightarrow y \text{ int. is } 12$$



3. a) i) $f(g(x))$: $f(x) = \sqrt{4x+3} \Rightarrow f(g(x)) = \sqrt{4(x^2-1)+3} = \sqrt{4x^2-1}$

(3)

$g(f(x))$: $g(x) = x^2 - 1 \Rightarrow g(f(x)) = (\sqrt{4x+3})^2 - 1 = 4x+2$

(ii) Domain $(f \circ g)(x)$ Step 1 Domain $g(x)$: $x \in \mathbb{R}$

(2)

Step 2 Domain of $f(g(x))$: $4x^2 - 1 \geq 0$ $\begin{cases} x \geq \frac{1}{2} \\ \text{OR} \\ x \leq -\frac{1}{2} \end{cases}$

$(g \circ f)(x)$

Step 1 Domain $f(x)$: $4x+3 \geq 0 \Rightarrow x \geq -\frac{3}{4}$

Step 2 Domain $g(f(x))$: $x \in \mathbb{R}$

4 a) x int

$$\frac{0}{1} = \frac{x^4 - 16}{3(x^2+1)(x^2-x-6)}$$

$$x^4 - 16 = 0$$

$$x^4 = 16$$

$$x = \pm 2$$

(2)

but $x = -2$ makes DEN = 0

$\Rightarrow x = 2$ only

b)

VA: let DEN = 0

$$3(x^2+1)(x^2-x-6) = 0$$

$$(x^2+1)(x^2-x-6) = 0$$

$$x^2+1=0$$

NO x

$$x^2-x-6=0$$

$$(x+2)(x-3)=0$$

$$x+2=0$$

$$x=-2$$

$$x-3=0$$

$$x=3$$

NUM = 0 when $x = -2$

$\Rightarrow x = 3$ is only VA.

y int

$$y = \frac{x^4 - 16}{3(x^2+1)(x^2-x-6)}$$

$$y = \frac{0^4 - 16}{3(0^2+1)(0^2-0-6)}$$

$$y = \frac{-16}{3(-6)} = \frac{-16}{-18} = \frac{8}{9}$$

c) HA:

(2) $\lim_{x \rightarrow \infty} f(x) = \frac{\infty}{\infty}$

$$\lim_{x \rightarrow \infty} \frac{\frac{x^4 - 16}{x^4}}{3\left(\frac{x^2+1}{x^2}\right)\left(\frac{x^2-x-6}{x^2}\right)}$$

$$\lim_{x \rightarrow \infty} \frac{1 - \frac{16}{x^4}}{3\left(1 - \frac{1}{x^2}\right)\left(1 - \frac{1}{x} - \frac{6}{x^2}\right)}$$

$$= \frac{1}{3} \Rightarrow \text{HA: } y = \frac{1}{3}$$

$$5a) \quad 9^x = 3^x + 12$$

$$3^{2x} - 3^x - 12 = 0$$

$$(3^x)^2 - 3^x - 12 = 0$$

$$(3^x - 4)(3^x + 3) = 0$$

$$3^x - 4 = 0 \quad 3^x + 3 = 0$$

$$3^x = 4 \quad 3^x = -3$$

$$\text{No } x$$

$$\ln 3^x = \ln 4$$

$$x \ln 3 = \ln 4$$

$$x = \frac{\ln 4}{\ln 3}$$

$$5b) \quad \log_2 x + \log_2 (x-14) = 5$$

$$\log_2 x(x-14) = 5$$

$$2^5 = x(x-14)$$

$$x^2 - 14x - 32 = 0$$

$$(x+2)(x-16) = 0$$

$x+2=0$	$x-16=0$
$x=-2$	$x=16$

DISCARD
domain
of $\log x$
is $x > 0$

$$6. a) y = \log_e(x+3) + 6$$

$$x = \log_e(y+3) + 6$$


$$x-6 = \log_e(y+3)$$

$$e^{x-6} = y+3$$

$$y = e^{x-6} - 3$$

$$\Rightarrow f^{-1}(x) = e^{x-6} - 3$$

b) Domain of $f(x)$: $x+3 > 0 \Rightarrow x \in \mathbb{R} \mid x > -3$
 Domain of $f^{-1}(x)$: (this will be Range of $f(x) \Rightarrow x \in \mathbb{R}$)



like $A^2 - A - 12 = 0$

$$(A-4)(A+3) = 0$$

Except we have

$$A = 3^x$$

$$5c) \quad 5^{\log_5 x^2} - 4 \cdot 3^{\log_3 (x+1)} = 8$$

$$x^2 - 4(x+1) = 8 \quad \otimes$$

$$x^2 - 4x - 4 - 8 = 0$$

$$x^2 - 4x - 12 = 0$$

$$(x+2)(x-6) = 0$$

$x+2=0$	$x-6=0$
$x=-2$	$x=6$

DISCARD
(domain)

\otimes let $(5^{\log_5 x^2}) = (A)$

$$\Rightarrow \log_5 A = \log_5 x^2$$

$$\Rightarrow A = x^2$$

$$\Rightarrow 5^{\log_5 x^2} = x^2$$

(Inverses of each other)

\Rightarrow Range $f(x)$ is $x \in \mathbb{R}$

Bonus:

a) Since Range of $f(x)$ is interval $(-2, 1)$

$$\Rightarrow -2 < f(x) < 1$$

$$\Rightarrow 0 \leq [f(x)]^2 < 4$$

① $[f(x)]^2$ cannot be Neg.

$$\textcircled{2} (-2)^2 = 4$$

\Rightarrow Range of $[f(x)]^2$ is interval $[0, 4)$

b) Since Range $f(x)$ is $(-2, 1)$

$\Rightarrow \exists x_1, x_2 (x_1 \neq x_2)$ and a

$$\text{Such that } f(x_1) = a \Rightarrow [f(x_1)]^2 = a^2$$

$$f(x_2) = -a \Rightarrow [f(x_2)]^2 = a^2$$

$\Rightarrow F(x) = [f(x)]^2$ is not one to one



$\Rightarrow F(x)$ is not Invertible

to have an Inverse a function must be 1 to 1.