




1. Find the area of the region enclosed by the curves  $y = x, y = x^3$ .


A.  $1/4$  B.  $-1/4$  C. 0 D.  $-1/8$  E. 1 F.  $1/2$  G.  $-1/2$  H. none of the above



$$A = \int_{-1}^1 |x^3 - x| dx = 2 \int_0^1 (x - x^3) dx = 2 \left( \frac{1}{2} - \frac{1}{4} \right) = \frac{1}{2}$$

2. Find the volume of the solid obtained by rotating the region bounded by the curves  $x = 2\sqrt{y}, x = 0, y = 5$  about the  $y$ -axis.

A.  $\pi/2$  B.  $50\pi$  C. 0 D.  $25\pi$  E.  $-50\pi$  F.  $\pi r^2$  G.  $-25\pi$  H. none of the above



$$V = \int_0^5 \pi r^2(y) dy = \int_0^5 4\pi y dy$$

$$= 2\pi \left[ y^2 \right]_{y=0}^{y=5} = 50\pi$$

3. Find the average value of the function  $g(t) = t/\sqrt{3+t^2}$  on the interval  $[1, 3]$ .

- A.  $2\sqrt{3}-1$  B.  $\sqrt{3}-2$  C. 1 D.  $\sqrt{3}-1$  E.  $2\sqrt{3}-2$  F.  $t^2/2$  G.  $1-\sqrt{3}$  H. none of the above

$$\bar{g} = \frac{I}{3-1} = \frac{1}{2} I \quad \text{with}$$

$$I = \int_1^3 \frac{t dt}{\sqrt{3+t^2}}$$


$$u = 3+t^2 \quad du = 2t dt$$

$$= \frac{1}{2} \int_4^{12} \frac{du}{\sqrt{u}} = u^{1/2} \Big|_{u=4}^{u=12} = \sqrt{12} - \sqrt{4} = 2(\sqrt{3}-1),$$

whence  $\bar{g} = \sqrt{3}-1$

4. Find the centroid of the region bounded by the curves  $y = 2 - x^2$ ,  $y = x$ .

- A.  $(-1, 2)$  B.  $(1/2, -2/5)$  C. 0 D.  $(-1/2, 1/2)$  E.  $(1/2, -1/2)$  F.  $1/2$  G.  $(-1/2, 2/5)$  H. none of the above



The area  $A = \int_{-2}^1 (2 - x^2 - x) dx$

$$= \left( 2x - \frac{x^3}{3} - \frac{x^2}{2} \right) \Big|_{x=-2}^{x=1}$$

$$= \left( 2 - \frac{1}{3} - \frac{1}{2} \right) - \left( -4 + \frac{8}{3} - 2 \right) = \frac{7}{6} + 6 - \frac{16}{6} = \frac{19}{2}$$

The coordinates of the centroid  $(\bar{x}, \bar{y})$  are

$$\bar{x} = \frac{I_x}{A}, \quad I_x = \int_{-2}^1 x(2 - x^2 - x) dx = \int_{-2}^1 (2x - x^3 - x^2) dx$$

$$= \left( x^2 - \frac{x^4}{4} - \frac{x^3}{3} \right) \Big|_{x=-2}^{x=1} = \left( 1 - \frac{1}{4} - \frac{1}{3} \right) - \left( 4 - 4 + \frac{8}{3} \right) = \frac{5}{12} - \frac{8}{3} = -\frac{9}{4} \Rightarrow \bar{x} = -\frac{1}{2}$$

$$\bar{y} = \frac{I_y}{A}, \quad I_y = \frac{1}{2} \int_{-2}^1 [(2-x^2)^2 - x^2] dx = \frac{1}{2} \int_{-2}^1 (x^4 - 5x^2 + 4) dx = \frac{1}{2} \left( \frac{x^5}{5} - \frac{5}{3}x^3 + 4x \right) \Big|_{x=-2}^{x=1}$$

$$= \frac{1}{2} \left( \frac{1}{5} - \frac{5}{3} + 4 + \frac{32}{5} - \frac{40}{3} + 8 \right) = \frac{1}{2} \left( \frac{33}{5} - \frac{45}{3} + 12 \right) = \frac{1}{2} \left( \frac{33}{5} - 3 \right) = \frac{9}{5} \Rightarrow \bar{y} = \frac{2}{5}$$

5. Which of the following improper integrals are convergent? (I)  $\int_0^{\infty} \frac{dx}{\sqrt{1+x}}$  (II)  $\int_0^1 \frac{dx}{x}$   
 (III)  $\int_{-\infty}^{\infty} xe^{-x^2} dx$

A. (I) only B. (II) only C. (III) only D. none E. (I) and (II) only F. all G. (II) and (III) only  
 H. none of the above

$$(I) \int_0^{\infty} \frac{dx}{\sqrt{1+x}} = 2(1+x)^{1/2} \Big|_{x=0}^{\infty} \text{ diverges}$$

$$(II) \int_0^1 \frac{dx}{x} = \ln x \Big|_{x=0}^{x=1} \text{ diverges}$$

$$(III) \int_{-\infty}^{\infty} xe^{-x^2} dx = \int_{-\infty}^0 xe^{-x^2} dx + \int_0^{\infty} xe^{-x^2} dx$$

$$= \frac{1}{2} \int_{-\infty}^0 e^{-x^2} d(x^2) + \frac{1}{2} \int_0^{\infty} e^{-x^2} d(x^2)$$

$$= \underbrace{-\frac{1}{2} \int_0^{\infty} e^{-t} dt}_{=-\frac{1}{2}} + \frac{1}{2} \int_0^{\infty} e^{-t} dt = \frac{1}{2} \text{ converges}$$

6. Determine whether the sequence  $a_n = n^2 \cos n / (1 + n^2)$  converges or diverges. If it converges, find its limit  $L$ .

A. diverges,  $L = 1$  B. converges,  $L = 1$  C. converges,  $L = \cos n$  D. diverges,  $L = \cos n$   
 E. converges,  $L = \pi$  F. converges,  $L = 0$  G. diverges H. none of the above

$$a_n = \frac{n^2 \cos n}{n^2 + 1} = \underbrace{\frac{n^2}{n^2 + 1}}_{\rightarrow 1} \cdot \cos n, \text{ so that}$$

the convergence of  $a_n$  is equivalent to the convergence of  $\cos n$

7. Which of the following series converge? (I)  $\sum_{k=2}^{\infty} \frac{k^2 \cos k}{k^4 - 1}$ , (II)  $\sum_{n=0}^{\infty} (-1)^{n-1} \frac{n^2}{n^3 + 1}$  (III)  $\sum_{n=1}^{\infty} \frac{n^2 - n}{3n^3 + 2}$

A. (I) only B. (II) only C. (III) only D. none  E. (I) and (II) only F. all G. (II) and (III) only H. none of the above

(I)  $\left| \frac{k^2 \cos k}{k^4 - 1} \right| \leq \frac{k^2}{k^4 - 1}$  which converges by the comparison test

(II)  $\left( \frac{x^2}{x^3 + 1} \right)' = \frac{2x(x^3 + 1) - 3x^4}{(x^3 + 1)^2} = \frac{2x - x^4}{(x^3 + 1)^2} < 0$  for  $x \geq 2$ , so that  $\frac{n^2}{n^3 + 1}$  is decreasing for  $n \geq 2$ , and  $\frac{n^2}{n^3 + 1} \rightarrow 0$ , whence the series converges

(III) diverges by the comparison with  $\sum_{n=1}^{\infty} \frac{1}{n}$  and using the limit comparison test

8. How many terms  $n$  of the series  $\sum_{n=1}^{\infty} \frac{1}{n^2}$  should one take so that the error  $R_n$  do not exceed 0.001?

A. 1999 B. 2001 C. 500 D. 501  E. 1000 F. 100 G. 2000 H. none of the above

Since  $f(x) = \frac{1}{x^2}$  is monotone decreasing,

$$R_n = \sum_{k=1}^{\infty} a_k - \sum_{k=1}^n a_k \leq \int_n^{\infty} \frac{dx}{x^2} = \left( -\frac{1}{x} \right) \Big|_{x=n}^{x=\infty} = \frac{1}{n},$$

whence  $R_n \leq 0.001$  if  $\frac{1}{n} \leq 0.001$

9. Find the convergence radius  $R$  and the convergence interval  $I$  of the power series  $\sum_{n=0}^{\infty} \frac{n}{2^n(n^2+1)} x^n$ .

- A.  $R = 1, I = [-1, 1]$     B.  $R = -1, I = [-2, 2]$     **C.  $R = 2, I = [-2, 2]$**     D.  $R = 2, I = (-2, 2]$   
 E.  $R = -2, I = (-2, 2)$     F.  $R = 1, I = [-1, 1]$     G.  $R = 1, I = (-1, 1]$     H. none of the above

For  $a_n = \frac{n x^n}{2^n (n^2 + 1)}$

$\left| \frac{a_{n+1}}{a_n} \right| = \frac{(n+1)(n^2+1)}{((n+1)^2+1)n} \cdot \left| \frac{x}{2} \right| \rightarrow \left| \frac{x}{2} \right|$ , so that the series converges for  $|x| < 2$ , diverges for  $|x| > 2$ , and  $R = 2$

at the endpoints:

for  $x = -2$   $a_n = (-1)^n \frac{n}{n^2+1}$ , and the series converges by the alternating test

for  $x = 2$   $a_n = \frac{n}{n^2+1}$ , and the series diverges by the comparison with the harmonic series

10. What is the third order entry  $a_3$  of the Taylor series of the function  $f(x) = \sqrt[3]{1-3x}$  at the point 0? By using the Taylor series find the limit  $L = \lim_{x \rightarrow 0} (\sqrt[3]{1-3x} - 1 + x)/x^2$ .

- A.  $a_3 = -\frac{5}{3}x^3, L = -1$**     B.  $a_3 = x^3, L = 1$     C.  $a_3 = x^3/2, L = -1/2$     D.  $a_3 = \frac{5}{3}x^3, L = 1$   
 E.  $a_3 = x^3, L = 0$     F.  $a_3 = x^3, L = -1$     G.  $a_3 = \frac{1}{2}x^3, L = 1$     H. none of the above

$$f(x) = (1-3x)^{1/3}$$

$$f(0) = 1$$

$$f'(x) = -(1-3x)^{-2/3}$$

$$f'(0) = -1$$

$$f''(x) = -2(1-3x)^{-5/3}$$

$$f''(0) = -2$$

$$f'''(x) = -10(1-3x)^{-8/3}$$

$$f'''(0) = -10$$

The Taylor series is  $1 - x - x^2 - \frac{5}{3}x^3 - \dots \Rightarrow a_3 = -\frac{5}{3}x^3$

$$\frac{f(x) - 1 + x}{x^2} = -1 - \frac{5}{3}x - \dots \Rightarrow L = -1$$

11. Find the solution of the differential equation  $y' = xe^y$  that satisfies the initial condition  $y(0) = 0$ .

- A.  $y = \ln(1 - x^2/2)$    B.  $y = 2 \ln x$    C.  $y = e^{-x}$    D.  $y = 1/x$    E.  $y = -2 \ln x$    F.  $y = e^{x^2-1}$   
 G.  $y = -\ln(1 - x^2/2)$    H. none of the above

$$\frac{dy}{dx} = xe^y \Rightarrow e^{-y} dy = x dx \Rightarrow \int e^{-y} dy = \int x dx + C$$

$$\Rightarrow -e^{-y} = \frac{x^2}{2} + C \quad ; \quad \text{for } x=0, y=0 \quad -1 = C$$

$$\Rightarrow e^{-y} = 1 - \frac{x^2}{2} \Rightarrow y = -\ln\left(1 - \frac{x^2}{2}\right)$$

12. Which of the following functions  $f = f(x, y)$  have the property that  $f_{xx} + f_{yy} = 0$ ?

(I)  $e^{x^2-y^2}$ , (II)  $x^3 + 3xy^2$ , (III)  $\ln(x^2 + y^2)$ , (IV)  $e^x \sin y - e^{-y} \cos x$ ?

- A. (II) and (IV)    B. (III) and (IV)   C. (I), (II) and (IV)   D. (II), (III) and (IV)   E. (I) and (III)   F. none  
 G. all   H. none of the above

(I)  $f_x = 2xe^{x^2-y^2}$ ,  $f_{xx} = 2e^{x^2-y^2} + 4x^2e^{x^2-y^2}$   
 $f_y = -2ye^{x^2-y^2}$ ,  $f_{yy} = -2e^{x^2-y^2} + 4y^2e^{x^2-y^2}$

(II)  $f_{xx} = 6x$ ,  $f_{yy} = 6x$

(III)  $f_x = \frac{2x}{x^2+y^2}$ ,  $f_{xx} = \frac{2(x^2+y^2) - 4x^2}{(x^2+y^2)^2} = \frac{-2x^2 + 2y^2}{(x^2+y^2)^2}$   
 Switching  $x$  and  $y$ ,  $f_{yy} = \frac{-2y^2 + 2x^2}{(x^2+y^2)^2}$

(IV)  $f_{xx} = e^x \sin y + e^{-y} \cos x$   
 $f_{yy} = -e^x \sin y - e^{-y} \cos x$

13. For  $z = x^2 - xy^2$  with  $x = 2s - t + u$  and  $y = st^2u^2$  find the partial derivatives  $\frac{\partial z}{\partial s}$ ,  $\frac{\partial z}{\partial t}$  and  $\frac{\partial z}{\partial u}$  at the point  $(s, t, u) = (2, 1, -1)$ .

- A. 0   B. (0, -8)   C. (1, 4, -4)   D.  $2x - 2xy$    E. (2, -1, 1)   **F. (-8, -32, 32)**  
 G.  $(2s - t + u)^2 - (2s - t + u)(st^2u^2)^2$    H. none of the above

$$\frac{\partial x}{\partial s} = 2, \quad \frac{\partial x}{\partial t} = -1, \quad \frac{\partial x}{\partial u} = 1$$

$$\frac{\partial y}{\partial s} = t^2u^2 = 1, \quad \frac{\partial y}{\partial t} = 2stu^2 = 4, \quad \frac{\partial y}{\partial u} = 2st^2u = -4$$

$$x(2, 1, -1) = 2, \quad y(2, 1, -1) = 2$$

$$\frac{\partial z}{\partial x} = 2x - y^2 = 0, \quad \frac{\partial z}{\partial y} = -2xy = -8$$

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial s} = -8$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial t} = -32$$

$$\frac{\partial z}{\partial u} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial u} = 32$$

14. Find the directional derivative of the function  $f(x, y, z) = x^2y - xyz$  at the point  $(1, -1, 2)$  in the direction of the vector  $\mathbf{v} = (-1, 2, -2)$ .

- A. 4   B. 4/3   C. 2/3   D. 0   **E. -4/3**   F. -2/3   G. -2   H. none of the above

$$\nabla f = (2xy - yz, x^2 - xz, -xy)$$

$$\Rightarrow \nabla f(1, -1, 2) = (0, -1, 1)$$

The length of vector  $\mathbf{v}$  is  $\sqrt{(-1)^2 + 2^2 + (-2)^2} = \sqrt{9} = 3$ ,  
 whence its normalization is  $\mathbf{u} = \mathbf{v}/3 = (-\frac{1}{3}, \frac{2}{3}, -\frac{2}{3})$ ,

so that

$$\begin{aligned} \mathcal{D}_{\mathbf{v}} f(1, -1, 2) &= \mathcal{D}_{\mathbf{u}} f(1, -1, 2) = (0, -1, 1) \cdot \left(-\frac{1}{3}, \frac{2}{3}, -\frac{2}{3}\right) \\ &= -\frac{4}{3} \end{aligned}$$



15. Find an equation of the tangent plane to the surface  $z = e^{x-y}$  at the point  $(2, 2, 1)$ .

- A.  $y = x + 1$    B.  $z = x + y$    C.  $z = x - y + 1$    D.  $x + y + z = 1$    E.  $x = y + z$    F.  $z = e^x - e^y$   
 G.  $z^2 = x - y$    H.  $z = x - y$  none of the above

$$z = f(x, y) \text{ with } f(x, y) = e^{x-y}$$

$$f_x(2, 2) = e^{x-y} \Big|_{\substack{x=2 \\ y=2}} = 1$$

$$f_y(2, 2) = -e^{x-y} \Big|_{\substack{x=2 \\ y=2}} = -1, \quad \text{whence the}$$

tangent plane equation is

$$\begin{aligned} z &= z_0 + f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0) \\ &= 1 + 1 \cdot (x - 2) + (-1)(y - 2) = x - y + 1 \end{aligned}$$