

Last Name: \_\_\_\_\_

First name: \_\_\_\_\_

Student Number: \_\_\_\_\_

# CHM 1311 A Final Exam Fall 2011

*Please keep your work covered at all times and keep your eyes on your own paper! Cheating or any appearance of cheating will result in an F in the course and possible expulsion from the university.*

There are 15 pages in this test. A periodic table, data tables, and a formula sheet are provided at the end. You may rip these pages off of the exam and use them to cover your work. Any scratch work should be done on the back of these pages.

Please show all work to receive partial credit.

You have 180 minutes to complete the exam.

## #1. (20 points) Short Answer Questions.

a) The *spdf* electron configuration of gallium is: \_\_\_\_\_

b) Which of the following ions is the strongest base?     $F^-$      $HO_2^-$      $N_3^-$      $NO_2^-$

c) Name the following compounds:

i.  $CrO_3$  \_\_\_\_\_

ii.  $CaSO_4 \cdot 4H_2O$  \_\_\_\_\_


d) The solubility of magnesium carbonate is highest in a buffer solution with a pH of

3.5                      8.5                      6.5                      10.5

e) For a second order reaction, the plot of \_\_\_\_\_ versus \_\_\_\_\_ will yield a straight line, where the slope is equal to \_\_\_\_\_.

f) A buffer solution is prepared from 0.450 M HCN and 0.450 M NaCN. The pH of this solution is \_\_\_\_\_.

g) In the following reaction, circle the oxidizing agent:



h) As the activation energy of a reaction increases, the reaction rate

INCREASES                      DECREASES                      STAYS CONSTANT


i) For an ideal gas, a graph of PV versus T will be a linear relationship with a slope that is directly proportional to the gas's:

KINETIC ENERGY                      MOLAR MASS                      ENTHALPY                      MASS

j) The concentration of lead (II) ions in a saturated solution of lead (II) chloride is \_\_\_\_\_.

k) A one litre balloon is filled with neon gas. A hole is made in the balloon and the gas effuses at a rate of 0.0106 mol/hr. If the same balloon is refilled with argon at the same pressure and temperature, its rate of effusion would be \_\_\_\_\_.

- l) Write the equilibrium constant expression for the following reaction, and choose the best means by which you could encourage the formation of hydrogen gas.





- i) Increase volume and add heat  
 ii) Add steam and remove heat  
 iii) Remove carbon monoxide and decrease volume  
 iv) Add hydrogen gas and increase volume

K =

- m) An orbital is a spherical region of space in which there is a high probability of finding an electron.                      TRUE                      FALSE

- n) The standard heat of formation of solid  $\text{Fe}(\text{OH})_3$  is  $-824 \text{ kJ/mol}$ . Write the chemical equation for the reaction to which this value applies.

- o) Which of the following would NOT produce a buffer? (HA = a weak acid)



- p) If the equilibrium constant for the reaction  $\text{HA} + \text{H}_2\text{O} \rightleftharpoons \text{H}_3\text{O}^+ + \text{A}^-$  is  $K_a$ , then the equilibrium constant for the reaction  $\text{A}^- + \text{H}_2\text{O} \rightleftharpoons \text{HA} + \text{OH}^-$  would be

$K_a$

$K_w/K_a$

$1/K_a$

$K_b$

- q) The maximum number of electrons with quantum numbers  $n = 2$  and  $\ell = 1$  is: \_\_\_\_\_.

**#2. (10 points)**

**(a) Draw a diagram of Bohr's model of the hydrogen atom showing the transition of an electron from the ground state to the  $n = 4$  level.**

**(b) What is the change in energy (in J) of this transition?**

**Answer :** \_\_\_\_\_

**(c) What is the significance of the sign in your answer for part (b)?**

**(d) Calculate the wavelength (in nm) that corresponds to this energy.**

**Answer :** \_\_\_\_\_

**#3. (10 points). You are given 200.0 mL of an aqueous solution that contains 0.250 M each of  $\text{Ag}^+$ ,  $\text{Pb}^{2+}$ , and  $\text{Ba}^{2+}$ . You wish to precipitate these cations sequentially by adding chromate to the solution.**

**(a) In what order will the solids precipitate as chromate is added to the above solution?**

**First = \_\_\_\_\_      Second = \_\_\_\_\_      Third = \_\_\_\_\_**

**(b) What concentration of chromate will cause the first precipitation?**

**Answer: \_\_\_\_\_**

**(c) At the point just before  $\text{BaCrO}_4$  begins to precipitate, what will be the concentration of all of the ions in solution?**

**$[\text{Ag}^+] =$  \_\_\_\_\_       $[\text{Pb}^{2+}] =$  \_\_\_\_\_       $[\text{Ba}^{2+}] =$  \_\_\_\_\_**

**#4. (10 points) Combustion of 1.110 g of a gaseous hydrocarbon yields 3.613 g of carbon dioxide and 1.109 g of water (and no other products).**

**a) What is the empirical formula of the hydrocarbon?**



**Answer:** \_\_\_\_\_

**b) A 0.288 g sample of the same hydrocarbon occupies a volume of 121 mL at 24.8°C and 753 mmHg. What is the molecular formula of the hydrocarbon?**

**Answer:** \_\_\_\_\_

**c) Write a balanced chemical equation for the combustion of the hydrocarbon.**

**#5. (10 points) In an environmental analysis for copper in soils, the copper must first be oxidized to copper(II) ions using permanganate ions in acidic solution, with the following half-reactions.**



**a) Determine the overall balanced redox reaction equation.**

**b) A 55.0 g sample of soil was found to react with 12.76 mL of a 0.280 M  $\text{KMnO}_4$  solution. What was the percent composition by mass of copper in the soil sample?**

**Answer:** \_\_\_\_\_

**#6. (10 points) You wish to prepare a buffer solution with pH = 9.45.**

- a) How many grams of  $(\text{NH}_4)_2\text{SO}_4$  would you add to 425 mL of 0.258 M  $\text{NH}_3$  to do this? You may assume that the solution's volume remains constant.  $K_b$  of  $\text{NH}_3 = 1.8 \times 10^{-5}$**

**Answer:** \_\_\_\_\_

- b) Which buffer component, and how much (in grams) would you add to 0.100 L of the buffer in part (a) to change its pH to 9.30? You may assume that the solution's volume remains constant.**

**Answer:** \_\_\_\_\_

**#7. (10 points) In the titration of 40.00 mL of 0.200 M HOCl by 0.500 M NaOH:**  
**(a) Calculate the initial pH (before any addition of NaOH).**

**Answer:** \_\_\_\_\_

**(b) What is the volume of NaOH needed to attain the equivalence point? Calculate the pH at the equivalence point.**

**Volume of NaOH:** \_\_\_\_\_

**pH:** \_\_\_\_\_

**#8. (10 points).** Steel is an alloy of iron and carbon, with iron being the major component. A steel ball bearing has a radius of 5.85 mm and a density of 7.75 g/cm<sup>3</sup>. If the ball bearing contains 0.25% carbon (by mass) and that the percent natural abundance of <sup>13</sup>C is 1.108%, how many <sup>13</sup>C atoms are present in the ball bearing?

*Recall: volume of a sphere =  $(4\pi r^3)/3$*

**Answer:** \_\_\_\_\_

**#9. (10 points)** A 0.3268 g sample of caffeine ( $\text{C}_8\text{H}_{10}\text{O}_2\text{N}_4$ , heat of combustion =  $-4.243 \times 10^3$  kJ/mol) undergoes complete combustion in a bomb calorimeter. The bomb calorimeter assembly has a heat capacity of 5.136 kJ/°C.

(a) What mass of oxygen is required for the complete combustion of the sample?



Answer: \_\_\_\_\_

(b) What will be the final temperature of the assembly if the initial temperature is 22.43°C?

Answer: \_\_\_\_\_

(c) What is the change in internal energy for the reaction?

Answer: \_\_\_\_\_

**Gas Laws**

$$PV = nRT$$

$$\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$$

$$P_T = P_1 + P_2 + P_3 + \dots$$

$$P_A = \chi_A P_T$$

$$d = \frac{m}{V} = \frac{P \cdot MM}{RT}$$

$$E_K = \frac{1}{2}mv^2$$

$$u_{rms} = \sqrt{\frac{3RT}{MM}}$$

$$\frac{\text{Rate } A}{\text{Rate } B} = \sqrt{\frac{MM_B}{MM_A}}$$

$$\left(P + \frac{n^2a}{V^2}\right)(V - nb) = nRT$$

**Equilibrium**

$$K_p = K_c(RT)^{\Delta n}$$

**Acid/Base**

$$pOH = -\log[OH^-]$$

$$pH = -\log[H^+]$$

$$pH + pOH = 14$$

$$K_a \times K_b = K_w$$

$$pH = pK_a + \log \frac{[A^-]}{[HA]}$$

$$pH = \frac{pK_{a1} + pK_{a2}}{2}$$

**Thermochemistry**

$$\Delta U = q + W$$

$$W_{\text{system}} = -P\Delta V = -\Delta nRT$$

$$\Delta H = \Delta U + P\Delta V$$

$$q_p = \Delta U + P\Delta V$$

$$q = ms\Delta T$$

$$q = n\Delta H$$

$$\Delta H_{\text{rxn}}^\circ = \sum n\Delta H_f^\circ(\text{pds}) - \sum n\Delta H_f^\circ(\text{rxts})$$

**The atom**

$$E = hv$$

$$c = v\lambda$$

$$E = -B/n^2$$

**Kinetics**

$$[A]_t = [A]_o - kt$$

$$\ln[A]_t = \ln[A]_o - kt$$

$$1/[A]_t = 1/[A]_o + kt$$

$$k = Ae^{(-E_a/RT)}$$

$$\ln(k_2/k_1) = (-E_a/R)(1/T_2 - 1/T_1)$$

**Other**

$$n = m/MM$$

$$C = n/V$$

$$\% \text{yield} = \frac{\text{actual yield}}{\text{theoretical yield}} \times 100\%$$

$$\chi_A = \frac{n_A}{n_T}$$

**Data For Water**

Density = 1.00 g/mL (at 25°C)

 $s = 2.13 \text{ J g}^{-1} \text{ K}^{-1}$  (solid) $s = 4.184 \text{ J g}^{-1} \text{ K}^{-1}$  (liquid) $s = 2.01 \text{ J g}^{-1} \text{ K}^{-1}$  (gas) $\Delta H^\circ_{\text{fus}} = 6.02 \text{ kJ mol}^{-1}$  $\Delta H^\circ_{\text{vap}} = 40.7 \text{ kJ mol}^{-1}$ **Constants and Conversion Factors**

1 mmHg = 1 torr    760 mmHg = 1 atm    1 atm = 101.325 kPa    1 atm = 1.013125 bar  
 1 cm<sup>3</sup> = 1 mL    1000 mL = 1 L    1000 L = 1 m<sup>3</sup>

Avogadro's Number	$N$	$6.022 \times 10^{23} \text{ mol}^{-1}$	
Boltzmann's constant	$k$	$1.30866 \times 10^{-23} \text{ J} \cdot \text{K}^{-1}$	
Faraday's constant	$F$	$96,485 \text{ C} \cdot \text{mol}^{-1}$	
Gas constant	$R$	$8.31451 \text{ J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$	
	$R$	$0.08206 \text{ atm} \cdot \text{L} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$	
	$R$	$8.31451 \text{ m}^3 \text{ Pa} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$	
	$R$	$0.0831451 \text{ bar} \cdot \text{L} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$	
Planck's constant	$h$	$6.62608 \times 10^{-34}$	J·s
Speed of Light	$c$	$2.99792458 \times 10^8$	m·s <sup>-1</sup>

**Table of Ionization Constants**


Acid			$K_a =$
Iodic acid	$\text{HIO}_3 + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{IO}_3^-$	$1.6 \times 10^{-1}$
Chlorous acid	$\text{HClO}_2 + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{ClO}_2^-$	$1.1 \times 10^{-2}$
Chloroacetic acid	$\text{HC}_2\text{H}_2\text{ClO}_2 + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{C}_2\text{H}_2\text{ClO}_2^-$	$1.4 \times 10^{-3}$
Nitrous acid	$\text{HNO}_2 + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{NO}_2^-$	$7.2 \times 10^{-4}$
Hydrofluoric acid	$\text{HF} + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{F}^-$	$6.6 \times 10^{-4}$
Formic acid	$\text{HCHO}_2 + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{CHO}_2^-$	$1.8 \times 10^{-4}$
Benzoic acid	$\text{HC}_7\text{H}_5\text{O}_2 + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{C}_7\text{H}_5\text{O}_2^-$	$6.3 \times 10^{-5}$
Hydrazoic acid	$\text{HN}_3 + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{N}_3^-$	$1.9 \times 10^{-5}$
Acetic acid	$\text{HC}_2\text{H}_3\text{O}_2 + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{C}_2\text{H}_3\text{O}_2^-$	$1.8 \times 10^{-5}$
Hypochlorous acid	$\text{HOCl} + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{OCl}^-$	$2.9 \times 10^{-8}$
Hydrocyanic acid	$\text{HCN} + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{CN}^-$	$6.2 \times 10^{-10}$
Phenol	$\text{HOC}_6\text{H}_5 + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{C}_6\text{H}_5\text{O}^-$	$1.0 \times 10^{-10}$
Hydrogen peroxide	$\text{H}_2\text{O}_2 + \text{H}_2\text{O} \rightleftharpoons$	$\text{H}_3\text{O}^+ + \text{HO}_2^-$	$1.8 \times 10^{-12}$

**Table of Solubility Product Constants**

Compound	$K_{\text{sp}}$	Compound	$K_{\text{sp}}$
Mg(OH) <sub>2</sub>	$1.2 \times 10^{-11}$	Ag <sub>2</sub> CrO <sub>4</sub>	$1.9 \times 10^{-12}$
AgCl	$1.8 \times 10^{-10}$	PbCrO <sub>4</sub>	$1.8 \times 10^{-14}$
CaSO <sub>4</sub>	$9.1 \times 10^{-6}$	BaCrO <sub>4</sub>	$2.1 \times 10^{-10}$
AgI	$1.5 \times 10^{-16}$	Hg <sub>2</sub> Cl <sub>2</sub>	$1.3 \times 10^{-18}$
PbI <sub>2</sub>	$8.7 \times 10^{-9}$	BaSO <sub>4</sub>	$1.1 \times 10^{-10}$
PbCl <sub>2</sub>	$1.9 \times 10^{-5}$	Ag <sub>2</sub> CO <sub>3</sub>	$8.5 \times 10^{-12}$

# Mokleur's Periodic table of the elements

																		18 VIIIA	
																		17 VIIA	
																		16 VIA	
																		15 VA	
																		14 IVA	
																		13 IIIA	
																		12 IIB	
																		11 IB	
																		10 VIII	
																		9 VII	
																		8 VIB	
																		7 VIIB	
																		6 VIB	
																		5 VB	
																		4 IVB	
																		3 IIIB	
																		2 IIA	
																		1 IA	
																		He 2 4.002602 Helium	
																		Ne 10 20.1797 Neon	
																		Ar 18 39.948 Argon	
																		Kr 36 83.80 Krypton	
																		Xe 54 131.29 Xenon	
																		Rn 86 222.0176 Radon	
																		Uuo 118 293 Ununoctium	
																		Fr 87 223.0197 Francium	
																		Ra 88 226.0254 Radium	
																		Ac 89 227.0278 Actinium	
																		Rf 104 261.11 Rutherfordium	
																		Db 105 262.11 Dubnium	
																		Sg 106 263.12 Seaborgium	
																		Bh 107 262.12 Bohrium	
																		Hs 108 264 Hassium	
																		Mt 109 266.1378 Meitnerium	
																		Uun 110 269 Ununium	
																		Uuu 111 272 Ununium	
																		Uuq 112 277 Ununium	
																		Uuh 116 289 Ununhexium	
																		Uuo 118 293 Ununoctium	
																		Lu 71 174.967 Lutetium	
																		Yb 70 173.04 Ytterbium	
																		Tm 69 168.93421 Thulium	
																		Er 68 167.26 Erbium	
																		Ho 67 164.93032 Holmium	
																		Dy 66 162.50 Dysprosium	
																		Ho 67 164.93032 Holmium	
																		Er 68 167.26 Erbium	
																		Tm 69 168.93421 Thulium	
																		Yb 70 173.04 Ytterbium	
																		Lu 71 174.967 Lutetium	
																		No 102 260.1053 Lawrencium	
																		Md 101 258.01 Mendelevium	
																		Fm 100 257.0951 Fermium	
																		Es 99 252.03 Einsteinium	
																		Cf 98 251.0796 Californium	
																		Bk 97 247.0703 Berkelium	
																		Cm 96 247 Curium	
																		Am 95 243.0614 Americium	
																		Pu 94 244.0642 Plutonium	
																		Np 93 237.04271 Neptunium	
																		U 92 238.02891 Uranium	
																		Pa 91 231.03588 Protactinium	
																		Th 90 232.03772 Thorium	
																		Pr 59 140.90765 Praseodymium	
																		Nd 60 144.242 Neodymium	
																		Pm 61 144.9127 Promethium	
																		Sm 62 150.36 Samarium	
																		Eu 63 151.9649 Europium	
																		Gd 64 157.25 Gadolinium	
																		Tb 65 168.93054 Terbium	
																		Dy 66 162.50 Dysprosium	
																		Ho 67 164.93032 Holmium	
																		Er 68 167.26 Erbium	
																		Tm 69 168.93421 Thulium	
																		Yb 70 173.04 Ytterbium	
																		Lu 71 174.967 Lutetium	



Under normal conditions, bold symbols correspond to solid state, bold italic correspond to liquid state, italic correspond to gaseous state and normal correspond to synthetic elements.